

On the extensions of the Diophantine triples in Gaussian integers

Adžaga, Nikola; Filipin, Alan; Franušić, Zrinka

Source / Izvornik: **Monatshefte für Mathematik, 2022, 197(4), 535 - 563**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:237:781222>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-12-02**

Repository / Repozitorij:

[Repository of the Faculty of Civil Engineering,
University of Zagreb](#)



ON THE EXTENSIONS OF THE DIOPHANTINE TRIPLES IN GAUSSIAN INTEGERS

NIKOLA ADŽAGA, ALAN FILIPIN, AND ZRINKA FRANUŠIĆ

ABSTRACT. A Diophantine m -tuple is a set of m distinct integers such that the product of any two distinct elements plus one is a perfect square. In this paper we study the extensibility of a Diophantine triple $\{k-1, k+1, 16k^3-4k\}$ in Gaussian integers $\mathbb{Z}[i]$ to a Diophantine quadruple. Similar one-parameter family, $\{k-1, k+1, 4k\}$, was studied in [9], where it was shown that the extension to a Diophantine quadruple is unique (with an element $16k^3-4k$). The family of the triples of the same form $\{k-1, k+1, 16k^3-4k\}$ was studied in rational integers in [6]. It appeared as a special case while solving the extensibility problem of Diophantine pair $\{k-1, k+1\}$, in which it was not possible to use the same method as in the other cases. As authors (Bugeaud, Dujella and Mignotte) point out, the difficulty appears because the gap between $k+1$ and $16k^3-4k$ is not sufficiently large. We find the same difficulty here while trying to use Diophantine approximations. Then we partially solve this problem by using linear forms in logarithms.

1. INTRODUCTION

A long-standing conjecture, motivated by work of Baker and Davenport [3], that there is no Diophantine quintuple, was proven by He, Togbé and Ziegler [11]. In other rings of integers, there are not many results. E.g. we find only about 10 papers solving similar problems in the ring of Gaussian integers. We can highlight [5] and [9], which deal with the extension of Diophantine triples from one-parameter families, and [1], which shows that there is no Diophantine m -tuple in imaginary quadratic number ring with $m \geq 43$.

We deal with a parametric family of triples $\{k-1, k+1, 16k^3-4k\}$, but we start with a general triple and show some results which are useful for any family of triples. Assume that a Diophantine triple $\{a, b, c\}$ in Gaussian integers $\mathbb{Z}[i]$ can be extended with a fourth element d . By eliminating d from the equations it satisfies ($ad+1=x^2$, $bd+1=y^2$ and $cd+1=z^2$), we get a system of two Pell-type equations with common unknown. We show that the structure of the solutions of this system is the same as in the rational integers case. A solution of this system gives us two simultaneous approximations of square roots close to 1. One can use Diophantine approximations in the general case (by assuming that $|c|$ is much bigger than $|b|$, say $|c| > |b|^{15}$), which was done in [1]. However, here we show that this is not useful for the triple of the form $\{k-1, k+1, 16k^3-4k\}$.

We also prove that the linear form in logarithms usually involved in approaching these problems is not zero under certain conditions. This might be useful in lowering the general upper bound, and we also use it here to partially resolve the extensibility problem of the triple $\{k-1, k+1, 16k^3-4k\}$.

Date: May 24, 2019.

2010 Mathematics Subject Classification. primary 11D09; secondary 11J68, 11J86.

Key words and phrases. Diophantine m -tuples, Diophantine approximation, Pell equations.

2. SYSTEM OF PELL-TYPE EQUATIONS

Let $\{a, b, c\} \subset \mathbb{Z}[i]$ be a Diophantine triple in Gaussian integers $\mathbb{Z}[i]$. Without loss of generality, we may assume $0 < |a| \leq |b| \leq |c|$. Then there are r, s and t in $\mathbb{Z}[i]$ such that $ab + 1 = r^2$, $ac + 1 = s^2$, $bc + 1 = t^2$. In [1], the following lemma was proven (for general imaginary quadratic number rings).

Lemma 2.1. *If $\{a, b, c\}$ is a Diophantine triple in the imaginary quadratic number ring $\mathbb{Z}[i]$ and $abc \neq 0$, then ab , ac and bc are not squares in $\mathbb{Z}[i]$.*

If there is $d \in \mathbb{Z}[i]$ such that $\{a, b, c, d\}$ is a Diophantine quadruple, then there are $x, y, z \in \mathbb{Z}[i]$ such that $ad + 1 = x^2$, $bd + 1 = y^2$, $cd + 1 = z^2$. Eliminating d implies that

$$(2.1) \quad az^2 - cx^2 = a - c$$

$$(2.2) \quad bz^2 - cy^2 = b - c.$$

These equations are similar to Pell's equations and their solutions have a very similar structure. The solutions of Pell-type equations ($x^2 - Dy^2 = N$) in imaginary quadratic rings are described in [8], as well as here, in a slightly different manner, adapted for the problem at hand.

Lemma 2.2. *There are positive integers i_0 and j_0 , elements $z_0^{(i)}, x_0^{(i)}, z_1^{(j)}, x_1^{(j)}$ of $\mathbb{Z}[i]$, for $i = 1, \dots, i_0$ and $j = 1, \dots, j_0$, such that:*

a) $(z_0^{(i)}, x_0^{(i)})$ are solutions of (2.1), while $(z_1^{(j)}, y_1^{(j)})$ are solutions of (2.2). The solutions denoted here are called fundamental.

b) Fundamental solutions satisfy the following inequalities:

$$1 \leq |x_0^{(i)}| \leq \sqrt{\frac{|a||c-a|}{|s|-1}}, \quad 1 \leq |z_0^{(i)}| \leq \sqrt{\frac{|c-a|}{|a|} + \frac{|c||c-a|}{|s|-1}},$$

$$1 \leq |y_1^{(j)}| \leq \sqrt{\frac{|b||c-b|}{|t|-1}}, \quad 1 \leq |z_1^{(j)}| \leq \sqrt{\frac{|c-b|}{|b|} + \frac{|c||c-b|}{|t|-1}}.$$

c) If (z, x) is the solution of (2.1), then there are $i \in \{1, \dots, i_0\}$ and $m \in \mathbb{Z}$ such that

$$z\sqrt{a} + x\sqrt{c} = (z_0^{(i)}\sqrt{a} + x_0^{(i)}\sqrt{c})(s + \sqrt{ac})^m.$$

If (z, y) is the solution of (2.2), then there are $j \in \{1, \dots, j_0\}$ and $n \in \mathbb{Z}$ such that

$$z\sqrt{b} + y\sqrt{c} = (z_1^{(j)}\sqrt{a} + y_1^{(j)}\sqrt{c})(t + \sqrt{bc})^n.$$

Proof. If (x, z) is the solution of (2.1), then the pairs $(x_m, y_m) \in \mathbb{Z}[i]^2$, defined as

$$(2.3) \quad x_m\sqrt{c} + z_m\sqrt{a} = (x\sqrt{c} + z\sqrt{a})(s + \sqrt{ac})^m$$

are also the solutions of (2.1) for every $m \in \mathbb{Z}$. We prove this inductively: for $m = 1$, we have $x_1\sqrt{c} + z_1\sqrt{a} = (x\sqrt{c} + z\sqrt{a})(s + \sqrt{ac}) = (sx + az)\sqrt{c} + (sz + cx)\sqrt{a}$, i. e. $x_1 = sx + az$, $z_1 = sz + cx$.

Let us note here that we have used Lemma 2.1. Then

$$\begin{aligned} az_1^2 - cx_1^2 &= a(sz + cx)^2 - c(sx + az)^2 = s^2(az^2 - cx^2) + ac(cx^2 - az^2) \\ &= s^2(a - c) + ac(c - a) = (s^2 - ac)(a - c) = a - c. \end{aligned}$$

Inductively it follows that (x_m, z_m) is the solution of (2.1) for every $m \in \mathbb{N}_0$. Analogously one resolves the case $m = -1$ to conclude that (x_m, z_m) is the solution (2.1) for every $m \in \mathbb{Z}$.

Let (x^*, z^*) be the solution such that $|x^*|$ is minimal among the solutions from the sequence $(x_m, z_m)_{m \in \mathbb{Z}}$ defined in (2.3). Let us denote the next and the previous solution in the sequence by (x', z') and (x'', z'') . More precisely, let $x' = sx^* + az^*$, $z' = sz^* + cx^*$ and $x'' = sx^* - az^*$, $z'' = sz^* - cx^*$. Then $|x'| \geq |x^*|$ and $|x''| \geq |x^*|$. On the other hand, by $|x'| + |x''| \geq |x' + x''| = 2|s||x^*|$ it follows that $|x'| \geq |sx^*|$ or $|x''| \geq |sx^*|$. In any case, we can conclude that the product $|x'x''| \geq |sx^*| \cdot |x^*|$. Since (z^*, x^*) is the solution of (2.1), we obtain equivalent inequalities $|(sx^* + az^*)(sx^* - az^*)| \geq |s| \cdot |x^*|^2$, $|(ac + 1)(x^*)^2 - a^2(z^*)^2| \geq |s| \cdot |x^*|^2$ and $|a(c - a) + (x^*)^2| \geq |s| \cdot |x^*|^2$.

We derive the upper bound on $|x^*|$ from the last inequality, $|a| \cdot |c - a| + |x^*|^2 \geq |a(c - a) + (x^*)^2| \geq |s| \cdot |x^*|^2$, so $|a| \cdot |c - a| \geq (|s| - 1)|x^*|^2$, and finally $|x^*|^2 \leq \frac{|a| \cdot |c - a|}{|s| - 1}$.

This bound on $|x^*|$ implies an upper bound on $|z^*|$, $|z^*| = \frac{|c(x^*)^2 - c + a|}{|a|} \leq \frac{|c||x^*|^2}{|a|} + \frac{|c - a|}{|a|}$. Without loss of generality, we may assume that $x_0 = x^*$, $z_0 = z^*$.

Analogously one gets the upper bounds on fundamental solutions of the equation (2.2). \square

From (c) part of the Lemma 2.2 one can obtain and solve the same recurrence relations as in the integer case (see [7]). More precisely, the following lemma holds

Lemma 2.3. *Every solution z of the equation (2.1) is contained in one of the following sequences*

$$(2.4) \quad v_0^{(i)} = z_0^{(i)}, \quad v_1^{(i)} = sz_0^{(i)} + cx_0^{(i)}, \quad v_{m+2}^{(i)} = 2sv_{m+1}^{(i)} - v_m^{(i)} \quad \text{for } i = 1, \dots, i_0.$$

Similarly, every solution z of the equation (2.2) is contained in one of the following sequences

$$(2.5) \quad w_0^{(j)} = z_1^{(j)}, \quad w_1^{(j)} = tz_1^{(j)} + cy_1^{(j)}, \quad w_{n+2}^{(j)} = 2tw_{n+1}^{(j)} - w_n^{(j)}, \quad \text{for } j = 1, \dots, j_0.$$

We skip the proof as it is the same as in the case of rational integers (see [7]).

If d extends the initial triple $\{a, b, c\}$, then z is the solution of both equations (2.1) and (2.2). Such z is contained in one of the sequences $v_m^{(i)}$ and in one $w_n^{(j)}$, i. e. $z = v_m^{(i)} = w_n^{(j)}$.

By solving the recurrences (2.4) and (2.5), we obtain

$$(2.6) \quad v_m^{(i)} = \frac{1}{2\sqrt{a}} \left((z_0^{(i)} \sqrt{a} + x_0^{(i)} \sqrt{c})(s + \sqrt{ac})^m + (z_0^{(i)} \sqrt{a} - x_0^{(i)} \sqrt{c})(s - \sqrt{ac})^m \right)$$

$$(2.7) \quad w_n^{(j)} = \frac{1}{2\sqrt{b}} \left((z_1^{(j)} \sqrt{b} + y_1^{(j)} \sqrt{c})(t + \sqrt{bc})^n + (z_1^{(j)} \sqrt{b} - y_1^{(j)} \sqrt{c})(t - \sqrt{bc})^n \right).$$

Let us denote $P = \frac{1}{\sqrt{a}}(z_0^{(i)} \sqrt{a} + x_0^{(i)} \sqrt{c})(s + \sqrt{ac})^m$, and $Q = \frac{1}{\sqrt{b}}(z_1^{(j)} \sqrt{b} + y_1^{(j)} \sqrt{c})(t + \sqrt{bc})^n$. By $z = v_m^{(i)} = w_n^{(j)}$, it follows that $P - \frac{c-a}{a}P^{-1} = Q - \frac{c-b}{b}Q^{-1}$.

Lemma 2.4. *If $|c| \geq 4|b|$ and $m, n \geq 3$, then $|P| > 12 \left| \frac{c}{a} \right|$ and $|Q| > 12 \left| \frac{c}{b} \right|$.*

Proof. Observe that $|s + \sqrt{ac}| \geq \sqrt{|ac|}$, since $\operatorname{Re} \sqrt{1 + \frac{1}{ac}} > 0 \Rightarrow \left| \sqrt{1 + \frac{1}{ac}} + 1 \right| > 1 \Rightarrow |\sqrt{ac} + 1 + \sqrt{ac}| > \sqrt{|ac|}$. Therefore

$$\begin{aligned} |P| &= \frac{1}{\sqrt{|a|}} |z_0 \sqrt{a} + x_0 \sqrt{c}| \cdot |s + \sqrt{ac}|^m \geq \frac{|c-a|}{\sqrt{|a|}} \frac{1}{|z_0 \sqrt{a} - x_0 \sqrt{c}|} \sqrt{|ac|}^m \\ &\geq \frac{|c-a|}{\sqrt{|a|}} \frac{1}{|x_0 \sqrt{c}| + |z_0| \sqrt{|a|}} |ac|^{3/2} \geq \frac{|c-|c|/4|}{\sqrt{|a|}} \frac{1}{3|c|/2 + |c| \sqrt{|a|}} |ac|^{3/2} \end{aligned}$$

Hence $P \geq \frac{3|a||c|^{3/2}}{6 + 4\sqrt{|a|}} > 12 \left| \frac{c}{a} \right|$. The last inequality is equivalent with $|a|^2|c|^{1/2} > 2(12 + 8\sqrt{|a|})$. Since $|c| > 4|a|$, it suffices to show that $2|a|^{5/2} > 2(12 + 8|a|^{1/2})$, which holds for $|a|^{1/2} \geq 2$.

Analogously, $|c| \geq 4|b|$ and $n \geq 3$ imply $|Q| > 12 \left| \frac{c}{b} \right|$. □

Therefore

$$\begin{aligned} ||P| - |Q|| &\leq |P - Q| = \left| \frac{c-a}{a}P^{-1} - \frac{c-b}{b}Q^{-1} \right| \leq \left| \frac{c}{a} - 1 \right| \frac{1}{|P|} + \left| \frac{c}{b} - 1 \right| \frac{1}{|Q|} \\ &\leq \left| \frac{c}{a} - 1 \right| \cdot \frac{1}{12} \left| \frac{a}{c} \right| + \left| \frac{c}{b} - 1 \right| \cdot \frac{1}{12} \left| \frac{b}{c} \right| \leq \frac{1}{12} \left(\left| 1 - \frac{a}{c} \right| + \left| 1 - \frac{b}{c} \right| \right) \leq \frac{5}{24}. \end{aligned}$$

It follows that $\left| \frac{|P| - |Q|}{|P|} \right| \leq \frac{5}{24} |P|^{-1} \leq \frac{5}{24} < 1$. We now apply the following simple Lemma B.2 from [13].

Lemma 2.5. *Let $\Delta > 0$ such that $|\Delta - 1| \leq a$. Then*

$$|\log \Delta| \leq \frac{-\log(1-a)}{a} |\Delta - 1|.$$

We obtain the following inequalities for $\Lambda = \frac{|Q|}{|P|}$:

$$\begin{aligned} |\log \Lambda| &= \left| \log \frac{|Q|}{|P|} \right| \leq \frac{24}{5} \log \frac{24}{19} \cdot \frac{5}{24} |P|^{-1} \leq \log \frac{24}{19} |P|^{-1} \leq \log \frac{24}{19} \sqrt{|a|} \frac{|x_0| \sqrt{|c|} + |z_0| \sqrt{|a|}}{|c-a|} |s + \sqrt{ac}|^{-m} \\ &\leq \log \frac{24}{19} \sqrt{|a|} \frac{3|c|^{3/2}/2 + |c|^{3/2}/2}{3|c|/4} |s + \sqrt{ac}|^{-m} = \frac{8}{3} \log \frac{24}{19} \sqrt{|ac|} |s + \sqrt{ac}|^{-m}. \end{aligned}$$

Lemma 2.6. *If $K = \frac{8}{3} \log \frac{24}{19}$, then*

$$|\log \Lambda| = \left| m \log |s + \sqrt{ac}| - n \log |t + \sqrt{bc}| + \log \frac{|\sqrt{b}(z_0 \sqrt{a} + x_0 \sqrt{c})|}{|\sqrt{a}(z_1 \sqrt{b} + y_1 \sqrt{c})|} \right| < K \sqrt{|ac|} |s + \sqrt{ac}|^{-m}.$$

3. LINEAR FORM IN LOGARITHMS IS NON-ZERO

Denote $\Lambda = \log \frac{|Q|}{|P|} = m \log |s + \sqrt{ac}| - n \log |t + \sqrt{bc}| + \log \frac{|\sqrt{b}(z_0^{(i)} \sqrt{a} + x_0^{(i)} \sqrt{c})|}{|\sqrt{a}(z_1^{(j)} \sqrt{b} + y_1^{(j)} \sqrt{c})|}$, so Λ is the

linear form in logarithms of algebraic numbers. This form is usually involved in solving the extension problems of Diophantine triples (for example, it was studied in [9] and [5]). The same linear form was also useful in solving some Thue equations [10]. It is usually shown that this form is not zero so that one can apply the famous Baker-Wüstholz theorem [4] and subsequently, bound the coefficients m and n . In rational integers, the proof that $\Lambda \neq 0$ is often trivial, but in quadratic fields it can cause considerable problems, as it happened in [9] and [10]. With some mild conditions, we prove that $\Lambda \neq 0$, and this is valid for an arbitrary imaginary quadratic field K and a, b, c in its ring of integers \mathcal{O}_K .

Lemma 3.1. *If $\{a, b, c\}$ is an extensible Diophantine triple, $\frac{|c|}{|a|} \notin \mathbb{Q}$ and $\frac{|c|}{|b|} \notin \mathbb{Q}$, then $\Lambda \neq 0$.*

Proof. The proof strategy is the same as in the proof of Lemma 5.2. in [9], but some details differ. Let us prove that, if $v_m = w_n$, then $|P| \neq |Q|$. $P \neq Q$ is easy, because otherwise $P^{-1} = Q^{-1}$, and then $v_m = w_n$ would imply $\frac{c-a}{a} = \frac{c-b}{b}$ and $b = a$ (since $c \neq 0$).

By definition, $P = A + B\alpha$, $Q = C + D\beta$, where $\alpha = \sqrt{\frac{c}{a}}$ and $\beta = \sqrt{\frac{c}{b}}$, $A, B, C, D \in K$. From $v_m = w_n$, we get $\frac{A+B\alpha+A-B\alpha}{2} = \frac{C+D\beta+C-D\beta}{2}$, ($v_m = A$, $w_n = C$) so $A = C$. Furthermore,

$$|P|^2 = (A + B\alpha)(\bar{A} + \bar{B}\bar{\alpha}) = A\bar{A} + \bar{A}B\alpha + A\bar{B}\bar{\alpha} + |B|^2|\alpha|^2 \text{ i. e.}$$

$$|P|^2 = p + u\alpha + \bar{u}\bar{\alpha} + q|\alpha|^2, \quad |Q|^2 = r + v\beta + \bar{v}\bar{\beta} + s|\beta|^2,$$

where $p, q, r, s \in \mathbb{Q}$ and $u, v \in K$. The idea is to show that the $1, \alpha, \bar{\alpha}, |\alpha|^2, \beta, \bar{\beta}$ and $|\beta|^2$ are linearly independent. We do this through three steps (claims **A**, **B** and **C**).

Claim A: The numbers $\alpha = \sqrt{\frac{c}{a}}, \beta = \sqrt{\frac{c}{b}}$ and $\sqrt{\frac{a}{b}}$ are algebraic of degree 2 over K .

Lemma 2.1 implies that $\frac{c}{a}, \frac{c}{b}$ and $\frac{a}{b}$ are not squares in K . □

Claim B: Basis for $K(\alpha, \bar{\alpha})$ over K is $B_\alpha = \{1, \alpha, \bar{\alpha}, |\alpha|^2\}$. Analogously, basis for $K(\beta, \bar{\beta})$ is $B_\beta = \{1, \beta, \bar{\beta}, |\beta|^2\}$.

If $\gamma \in K(\alpha, \bar{\alpha})$, then $\gamma = \sum q_{ij}\alpha^i\bar{\alpha}^j$, where $q_{ij} \in K$. However, since $\alpha^2 = \frac{c}{a}$ and $\bar{\alpha}^2$ are in K and $\alpha\bar{\alpha} = |\alpha|^2$, it follows that one can write γ as $\gamma = q_0 + q_1\alpha + q_2\bar{\alpha} + q_3|\alpha|^2$.

To prove that B_α is linearly independent set over K , we first show that $\{1, \alpha, \bar{\alpha}\}$ is linearly independent. Assume the contrary. Then $\bar{\alpha} = A + B\alpha$ for $A, B \in K$. This implies $\bar{\alpha}^2 - A^2 - B^2\alpha^2 = 2AB\alpha$.

Hence, if $AB \neq 0$, then $\alpha = \frac{\bar{\alpha}^2 - A^2 - B^2\alpha^2}{2AB} \in K$, which contradicts the claim **A**. If $B = 0$, then

$\bar{\alpha} = A \in K$. This would imply that $\alpha = \sqrt{\frac{c}{a}}$ is in K , i. e. $\frac{c}{a} = \frac{x^2}{y^2}$ for some $x, y \in \mathcal{O}_K$, so $\frac{|c|}{|a|} = \frac{|x|^2}{|y|^2} \in \mathbb{Q}$, which contradicts the lemma hypothesis.

If $A = 0$, then $\bar{\alpha} = B\alpha$, so again $\frac{|c|}{|a|} = |\alpha|^2 = \alpha\bar{\alpha} = B\alpha^2 \in K \cap \mathbb{R}$, i. e. again it follows that $\frac{|c|}{|a|} \in \mathbb{Q}$. Therefore, $\{1, \alpha, \bar{\alpha}\}$ is linearly independent set over K .

For B_α , it suffices to show that there are no $A, B, C \in K$ such that

$$(3.1) \quad |\alpha|^2 = A + B\alpha + C\bar{\alpha}.$$

We first prove $C \neq 0$. The contrary would imply $|\alpha|^4 = A^2 + B^2\alpha^2 + 2AB\alpha$ and $2AB\alpha \in K$. Since $\alpha \notin K$, it follows that $AB = 0$. If $B = 0$, then $|\alpha|^2 = A \in \mathbb{Q}$, which contradicts the lemma hypothesis. If $A = 0$, then $|\alpha|^2 = B\alpha$, so $\bar{\alpha} = B \in K$, which contradicts the claim **A**. Therefore, $C \neq 0$.

By multiplying (3.1) by α , we get $\alpha^2\bar{\alpha} = A\alpha + B\alpha^2 + C|\alpha|^2$, which implies $C|\alpha|^2 = -A\alpha - B\alpha^2 + \alpha^2\bar{\alpha}$, i. e. $|\alpha|^2 = -\frac{B}{C}\alpha^2 - \frac{A}{C}\alpha + \frac{1}{C}\alpha^2\bar{\alpha}$. Since $\{1, \alpha, \bar{\alpha}\}$ is linearly independent, the last obtained equality together with (3.1) implies that $A = -\frac{B}{C}\alpha^2$, $B = -\frac{A}{C}$, $C = \frac{1}{C}\alpha^2$. This implies $C^2 = \alpha^2$, which contradicts the claim **A** (α^2 is not a square in K). \square

Claim C: The set $B = \{1, \alpha, \bar{\alpha}, |\alpha|^2, \beta, \bar{\beta}, |\beta|^2\}$ is linearly independent over K .

First we show that $\beta, \bar{\beta}$ and $|\beta|^2$ are not in $K(\alpha, \bar{\alpha})$. Let us assume that β can be written as

$$(3.2) \quad \beta = A + B\alpha + C\bar{\alpha} + D|\alpha|^2,$$

for some $A, B, C, D \in K$. Then

$$\beta^2 = A^2 + B^2\alpha^2 + C^2\bar{\alpha}^2 + D^2|\alpha|^4 + 2AB\alpha + 2AC\bar{\alpha} + 2AD|\alpha|^2 + 2BC|\alpha|^2 + 2BD\alpha^2\bar{\alpha} + 2CD\bar{\alpha}^2\alpha.$$

By $\beta^2 \in K$, it follows that the coefficients of algebraic numbers $\alpha, \bar{\alpha}$ and $|\alpha|^2$ are zero, i. e.

$$(3.3) \quad AB + CD\bar{\alpha}^2 = 0,$$

$$(3.4) \quad AC + BD\alpha^2 = 0,$$

$$(3.5) \quad AD + BC = 0.$$

By (3.3) and (3.5), multiplying (3.3) by A or C , it follows that $A^2 = C^2\bar{\alpha}^2$ or $B = D = 0$. Similarly, from (3.4) and (3.5), it follows that $A^2 = B^2\alpha^2$ or $C = D = 0$, while (3.3) and (3.4) imply that $A^2 = D^2|\alpha|^4$ or $B = C = 0$. There are four cases now.

- $B = C = D = 0$. By (3.2), it follows that $\beta \in K$, which contradicts the claim **A**.
- $A = B = C = 0$. Then $\beta = D|\alpha|^2$ and $\frac{|c|}{|b|} = |D|^2 \frac{|c|^2}{|a|^2} \in \mathbb{Q}$, which contradicts the lemma hypothesis.
- $B \neq 0, C = D = 0$. Hence $\beta = B\alpha$, i. e. $\sqrt{\frac{a}{b}} = B \in K$, which contradicts the claim **A**.
- $B \neq 0$ and at least one of C and D is non-zero. Then $A^2 = C^2\bar{\alpha}^2 = B^2\alpha^2 = D^2|\alpha|^4$, so $\beta^2 = 4A^2$, which again contradicts **A**.

Therefore, β cannot be written as a linear combination of elements in B_α . The same holds for $\bar{\beta}$ and $|\beta|^2$ and is proven identically.

The set $L[\{1, \alpha, \bar{\alpha}, |\alpha|^2\}]$ (spanned by B_α) is closed on inversion. Namely,

$$\frac{1}{A + B\alpha + C\bar{\alpha} + D|\alpha|^2} = \frac{(A + B\alpha) - (C\bar{\alpha} + D|\alpha|^2)}{K + L\alpha} = \frac{((A + B\alpha) - (C\bar{\alpha} + D|\alpha|^2))(K - L\alpha)}{K^2 - L^2\alpha^2},$$

where $K = A^2 + B^2\alpha^2 - C^2\bar{\alpha}^2 - D^2|\alpha|^4$, $L = 2(AB - CD\bar{\alpha}^2)$.

Now we show that $\bar{\beta}$ cannot be written as linear combination of elements in $B_\alpha \cup \{\beta\}$. Analogously one shows the linear independence of sets $B_\alpha \cup \{\bar{\beta}, |\beta|^2\}$ and $B_\alpha \cup \{\beta, |\beta|^2\}$. Namely, that implies $\bar{\beta} = q_1 + q_2\alpha + q_3\bar{\alpha} + q_4|\alpha|^2 + q_5\beta$ and $q_5 \neq 0$. Therefore

$$\begin{aligned} \bar{\beta}^2 = & q_1^2 + q_2^2\alpha^2 + q_3^2\bar{\alpha}^2 + q_4|\alpha|^4 + q_5^2 + 2(q_1q_2 + q_3q_4\bar{\alpha}^2)\alpha + 2(q_1q_3 + q_2q_4\alpha^2)\bar{\alpha} + 2(q_1q_4 + q_2q_3)|\alpha|^2 + \\ & + 2q_1q_5\beta + 2(q_2q_5\alpha + q_3q_5\bar{\alpha})\beta + 2q_4q_5|\alpha|^2\beta, \end{aligned}$$

so we see that $2q_5\beta(q_1 + q_2\alpha + q_3\bar{\alpha} + q_4|\alpha|^2) \in L[\{1, \alpha, \bar{\alpha}, |\alpha|^2\}]$. By $q_5 \neq 0$, it follows that $q_1 + q_2\alpha + q_3\bar{\alpha} + q_4|\alpha|^2 = 0$, i. e. $q_1 = q_2 = q_3 = q_4 = 0$. However, that means $\bar{\beta} = q_5\beta$ for some $q_5 \in K$, which implies $|\beta|^2 = q_5\beta^2 \in K \cap \mathbb{R}$, i. e. $|\beta|^2 \in \mathbb{Q}$, contradicting the lemma hypothesis.

We get the contradiction in a similar way if we assume that $|\beta|^2$ can be written as linear combination of elements in $\{1, \alpha, \bar{\alpha}, |\alpha|^2, \beta, \bar{\beta}\}$. By $|\beta|^2 = q_1 + q_2\alpha + q_3\bar{\alpha} + q_4|\alpha|^2 + q_5\beta + q_6\bar{\beta}$, it follows that $2(q_1 + q_2\alpha + q_3\bar{\alpha} + q_4|\alpha|^2 + q_5q_6)(q_5\beta + q_6\bar{\beta}) \in L[\{1, \alpha, \bar{\alpha}, |\alpha|^2\}]$, so $q_5\beta + q_6\bar{\beta} = 0$, which would again imply $|\beta|^2 \in \mathbb{Q}$ or $q_1 + q_2\alpha + q_3\bar{\alpha} + q_4|\alpha|^2 + q_5q_6 = 0$. Since B_α is linearly independent, it follows that $q_2 = q_3 = q_4 = 0$ and $q_1 + q_5q_6 = 0$. Hence $|\beta|^2 = q_1 + q_5\beta + q_6\bar{\beta}$, but this contradicts the linear independence of B_β . \square

Let us remind the reader that, prior to these three claims, we have shown that, from $|P|^2 = (A + B\alpha)(\bar{A} + \bar{B}\bar{\alpha})$ and a similar equality for $|Q|^2$, it follows that

$$|P|^2 = p + u\alpha + \bar{u}\bar{\alpha} + q|\alpha|^2, \quad |Q|^2 = r + v\beta + \bar{v}\bar{\beta} + s|\beta|^2,$$

where $p, q, r, s \in \mathbb{Q}$, and $u, v \in K$. Since we want to prove that $|P| \neq |Q|$, it suffices to show that $|P|^2 \neq |Q|^2$. If $|P|^2 = |Q|^2$, this would imply $(p - r) + u\alpha + \bar{u}\bar{\alpha} + q|\alpha|^2 - v\beta - \bar{v}\bar{\beta} - s|\beta|^2 = 0$, so the claim **C** implies that $p - r = u = q = v = s = 0$, i. e. $P = A = C = Q$, which we have already proven to be impossible. Therefore $|P| \neq |Q|$, which implies that $\Lambda = \log \frac{|P|}{|Q|} \neq 0$. \square

The statement of this lemma depends on the system of equations chosen at the beginning. However, one easily sees that the analogous claim holds even if one begins with a different system (e. g. $az^2 - cx^2 = a - c$, $ay^2 - bx^2 = a - b$). Choosing which system to deal with usually depends on being able to find all the fundamental solutions for one of the equations. Regardless of which system is chosen, one can use this lemma.

4. SYSTEM OF PELL-TYPE EQUATIONS FOR TRIPLES OF THE FORM $\{k - 1, k + 1, 16k^3 - 4k\}$

The set $\{k - 1, k + 1, 16k^3 - 4k\}$ is a Diophantine triple for every Gaussian integer k . Denote by $s = 4k^2 - 2k - 1$, $t = 4k^2 + 2k - 1$ (so $(k - 1)(16k^3 - 4k) + 1 = s^2$ and $(k + 1)(16k^3 - 4k) + 1 = t^2$). Assume now that d extends this Diophantine triple, i. e., that $\{k - 1, k + 1, 16k^3 - 4k, d\}$ is a Diophantine

quadruple in $\mathbb{Z}[i]$. There exist $x, y, z \in \mathbb{Z}[i]$ such that

$$(k-1)d+1 = x^2, (k+1)d+1 = y^2, (16k^3-4k)d+1 = z^2.$$

By eliminating d , we obtain the system

$$(4.1) \quad (k+1)x^2 - (k-1)y^2 = 2,$$

$$(4.2) \quad (16k^3-4k)x^2 - (k-1)z^2 = 16k^3-5k+1$$

Let $|k| > 3$. By Lemma 2.4 of [9], all the solutions of the equation (4.1) are given by $x = \pm V_n$, where (V_n) is a recurrent sequence defined by

$$(4.3) \quad V_0 = 1, V_1 = 2k-1, V_{n+2} = 2kV_{n+1} - V_n, \text{ for all } n \geq 0.$$

All solutions of the equation (4.2) are described in the following lemma, which follows from Lemma 2.2.

Lemma 4.1. *Let $k \in \mathbb{Z}[i] \setminus \{0, 1\}$. There are $j_0 \in \mathbb{N}$, $x_1^{(j)}, z_1^{(j)} \in \mathbb{Z}[i]$, $j = 1, \dots, j_0$ such that*

- a) $(x_1^{(j)}, z_1^{(j)})$ is the solution of (4.2) for all $j = 1, \dots, j_0$,
- b) these fundamental solutions are bounded as follows:

$$|x_1^{(j)}|^2 \leq \frac{|16k^3-5k+1||k-1|}{|4k^2-2k-1|-1}$$

$$|z_1^{(j)}|^2 \leq \frac{|16k^3-4k||16k^3-5k+1|}{|4k^2-2k-1|-1} + \frac{|16k^3-5k+1|}{|k-1|}$$

for all $j = 1, \dots, j_0$,

- c) for each solution (x, z) of (4.2) there are $j \in \{1, \dots, j_0\}$ and $m \in \mathbb{Z}$ such that

$$x\sqrt{16k^3-4k} + z\sqrt{k-1} = (x_1^{(j)}\sqrt{16k^3-4k} + z_1^{(j)}\sqrt{k-1}) \cdot (4k^2-2k-1 + \sqrt{(k-1)(16k^3-4k)})^m.$$

Hence, the solution x of the equation (4.2) is $x = \pm W_m^{(j)}$ for some $j \in \{1, \dots, j_0\}$ and $m \in \mathbb{N}_0$, where the sequence $(W_m^{(j)})_m$ is recurrently defined by

$$W_0^{(j)} = x_1^{(j)}, W_1^{(j)} = x_1^{(j)}(4k^2-2k-1) + z_1^{(j)}(k-1), W_{m+2}^{(j)} = 2(4k^2-2k-1)W_{m+1}^{(j)} - W_m^{(j)}, m \geq 0.$$

For the time being, we omit the upper index (j) .

If x is the solution of both (4.1) and (4.2), then $x = V_n = W_m$. We are looking for the common elements of the sequences $(V_n)_n$ and $(W_m)_m$.

We apply the congruence method now. Observe the remainders that (V_n) and (W_m) leave when divided by $s = 4k^2 - 2k - 1$. The following lemma is easily proven by induction.

Lemma 4.2. *For the sequences $(V_n)_n$ and $(W_m)_m$ it holds*

$$V_n \equiv 0, \pm 1, \pm(2k-1) \pmod{4k^2-2k-1} \text{ and}$$

$$W_m \equiv \pm x_1, \pm z_1(k-1) \pmod{4k^2-2k-1}$$

for all indices n and m .

By analysing these combinations we can conclude that, when $|k| > 17$, all fundamental solutions (x_1, z_1) which generate sequences $(W_m)_m$ that can intersect the sequence $(V_n)_n$, are given by the set

$$(x_1, z_1) \in \{(\pm 1, \pm 1), (\pm k, \pm(4k^2 + 2k - 1)), (\pm(2k - 1), \pm(8k^2 - 1))\}.$$

E. g. if $x_1 \equiv 0 \pmod{4k^2 - 2k - 1}$, then $x_1 = 0$ or $|x_1| \geq 4|k|^2 - 2|k| - 1$. However, in the latter case, the bound given in Lemma 4.1 implies that

$$(4|k|^2 - 2|k| - 1)^2 \leq \frac{|16k^3 - 5k + 1||k - 1|}{|4k^2 - 2k - 1| - 1},$$

i. e. $(4|k|^2 - 2|k| - 1)^2(|4k^2 - 2k - 1| - 1) \leq |16k^3 - 5k + 1||k - 1|$, which implies $16|k|^4 + 16|k|^3 + 5|k|^2 + 4|k| + 1 \geq 64|k|^6 - 96|k|^5 - 16|k|^4 - 56|k|^3 - 4|k|^2 - 10|k| - 2$, and is in turn equivalent to $-64|k|^6 + 96|k|^5 + 32|k|^4 + 72|k|^3 + 9|k|^2 + 14|k| + 3 \geq 0$, which is obviously impossible for large $|k|$ (one can determine that the largest zero of the left-hand side polynomial in $|k|$ is approximately 2.04414). For $x_1 \equiv \pm 1, \pm(2k - 1)$, we similarly exclude all the possibilities except $x_1 = \pm 1$ and $x_1 = \pm(2k - 1)$, which gives us the solutions $(\pm 1, \pm 1), (\pm(2k - 1), \pm(8k^2 - 1))$.

If $z_1(k - 1) \equiv 0, \pm 1, \pm(2k - 1) \pmod{4k^2 - 2k - 1}$, then $z_1 = u(4k^2 - 2k - 1) + r$ where u is a Gaussian integer, while $r \in \{0, \pm 4k, \pm(4k + 2)\}$, since $-4k - 2$ is the multiplicative inverse of $k - 1$ modulo $4k^2 - 2k - 1$. The equation (4.2) implies $k \mid 1 - z_1^2$. On the other hand, $z_1 \equiv -u + r \equiv -u, -u \pm 2 \pmod{k}$, so $k \mid 1 - u^2$ or $k \mid 1 - (u \pm 2)^2$. If $|u| \leq 2$, then $|1 - u^2| \leq 1 + |u|^2 \leq 5$ and $|1 - (u \pm 2)^2| \leq |u|^2 + 4|u| + 5 \leq 17$, and the obtained divisibility cannot hold if $|k| > 17$, except when $u = \pm 1$. Here we get the solutions $(\pm k, \pm(4k^2 + 2k - 1))$ for $u = \pm 1$ and $z_1 = u(4k^2 - 2k - 1)$. It is not possible that both $u = \pm 1$ and $r = \mp(4k + 2)$ hold, because then from the equation (4.2) it follows that $x_1^2 = \frac{(k-1)(4k^2-6k-3)^2+(16k^3-5k+1)}{16k^3-4k} \in \mathbb{Z}[i]$. This implies $2k^2 - k \mid 8k + 8$, which is impossible for $|k| > 17$ (since $8k + 8 = 0$ or $8|k| + 8 \geq 2|k|^2 - |k|$). If $|u| \geq \sqrt{5}$, repeating the juxtaposition with upper bound from Lemma 4.1, we see that this cannot hold for $|k| > 17$: $|z_1| \geq \sqrt{5}(4|k|^2 - 2|k| - 1) - 4|k| - 2 = 4\sqrt{5}|k|^2 - (4 + 2\sqrt{5})|k| - (2 + \sqrt{5})$, so Lemma 4.1 implies

$$\begin{aligned} 80|k|^4 - 32\sqrt{5}|k|^3 - 80|k|^3 - 4|k|^2 + 16\sqrt{5}|k| + 36|k| + 4\sqrt{5} + 9 &\leq \\ &\leq \frac{(16|k|^3 + 4|k|)(16|k|^3 + 5|k| + 1)}{4|k|^2 - 2|k| - 2} + \frac{16|k|^3 + 5|k| + 1}{|k| - 1}. \end{aligned}$$

It follows that $-64|k|^7 + (544 + 128\sqrt{5})|k|^6 + (-256 - 192\sqrt{5})|k|^5 + (-488 - 64\sqrt{5})|k|^4 + (332 + 144\sqrt{5})|k|^3 + (40 + 24\sqrt{5})|k|^2 + (-88 - 32\sqrt{5})|k| - 8\sqrt{5} - 20 \leq 0$, which does not hold for $|k| > 12.019$.

This proves the following lemma.

Lemma 4.3. *If $|k| > 17$ and the system of equations (4.1) and (4.2) has a solution $x \neq \pm 1$, then there are positive integers m and n , and $1 \leq j \leq 6$, such that*

$$V_n = \pm W_m^{(j)},$$

where the sequences $(W_m^{(j)})$ are given with the following initial conditions

$$\begin{aligned} W_0^{(1)} &= 1, & W_1^{(1)} &= 4k^2 - k - 2, \\ W_0^{(2)} &= 1, & W_1^{(2)} &= 4k^2 - 3k, \\ W_0^{(3)} &= k, & W_1^{(3)} &= 8k^3 - 4k^2 - 4k + 1, \\ W_0^{(4)} &= k, & W_1^{(4)} &= 2k - 1, \\ W_0^{(5)} &= 2k - 1, & W_1^{(5)} &= 16k^3 - 16k^2 - k + 2, \\ W_0^{(6)} &= 2k - 1, & W_1^{(6)} &= k, \end{aligned}$$

and all the other elements are defined by

$$W_{m+2}^{(j)} = 2(4k^2 - 2k - 1)W_{m+1}^{(j)} - W_m^{(j)}, \quad m \geq 0.$$

Observe that the sequences $(W_m^{(j)})_{j=1,\dots,6}$ intersect with $(V_n)_n$ at $\{1\}$, $\{1\}$, $\{8k^3 - 4k^2 - 4k + 1\}$, $\{2k - 1\}$, $\{2k - 1\}$, $\{2k - 1, 8k^3 - 4k^2 - 4k + 1\}$, respectively. These intersections correspond to the extensions $d \in \{0, 4k, 64k^5 - 48k^3 + 8k\}$.

5. LOWER BOUND FOR THE SOLUTIONS

With the aim of obtaining a lower bound for the solution $|x|$, we determine the remainders of the elements of sequences from Lemma 4.3 modulo $4k(k - 1)$. In this section, unless otherwise specified, we assume that $|k| > 17$. By calculating the first few elements of the sequences, we get

$$\begin{aligned} (V_n)_{n \geq 0} &\equiv (1, 2k - 1, 2k - 1, 1, 1, 2k - 1, \dots) && \pmod{4k(k - 1)}, \\ (W_m^{(1)})_{m \geq 0} &\equiv (1, 3k - 2, -2k + 3, 5k - 4, -4k + 5, 7k - 6, \dots) && \pmod{4k(k - 1)}, \\ (W_m^{(2)})_{m \geq 0} &\equiv (1, k, 2k - 1, -k + 2, 4k - 3, -3k + 4, \dots) && \pmod{4k(k - 1)}, \\ (W_m^{(3)})_{m \geq 0} &\equiv (k, 1, 3k - 2, -2k + 3, 5k - 4, -4k + 5, \dots) && \pmod{4k(k - 1)}, \\ (W_m^{(4)})_{m \geq 0} &\equiv (k, 2k - 1, -k + 2, 4k - 3, -3k + 4, 6k - 5, \dots) && \pmod{4k(k - 1)}, \\ (W_m^{(5)})_{m \geq 0} &\equiv (2k - 1, -k + 2, 4k - 3, -3k + 4, 6k - 5, \dots) && \pmod{4k(k - 1)}, \\ (W_m^{(6)})_{m \geq 0} &\equiv (2k - 1, k, 1, 3k - 2, -2k + 3, 5k - 4, \dots) && \pmod{4k(k - 1)}. \end{aligned}$$

Lemma 5.1. *Let k be a Gaussian integer (of absolute value greater than 1). For the sequence $(V_n)_n$ defined in (4.3), it holds that $V_n \equiv 1 \pmod{4k(k-1)}$ if $n \equiv 0, 3 \pmod{4}$, while $V_n \equiv 2k-1 \pmod{4k(k-1)}$ for $n \equiv 1, 2 \pmod{4}$.*

For the sequence $(W_m^{(1)})_m$ defined in Lemma 4.3, its elements $W_{2m}^{(1)} \equiv -2mk + 2m + 1 \pmod{4k(k-1)}$ and $W_{2m+1}^{(1)} \equiv (2m + 3)k - 2m - 2 \pmod{4k(k-1)}$ for all $m \in \mathbb{N}_0$. This sequence $(W_m^{(1)})_m$ is increasing in absolute value, and the inequality $|W_m^{(1)}| \geq (8|k|^2 - 4|k| - 3)^{m-1}$ holds for all $m \geq 0$. Similarly, the other sequences $(W_m^{(j)})_m$, for $j = 2, \dots, 6$, are increasing (in absolute value) after the index $m = 1$ and

$|W_m^{(j)}| \geq (8|k|^2 - 4|k| - 3)^{m-1}$. The following congruences also hold

$$\begin{aligned}
W_{2m}^{(2)} &\equiv 2mk - (2m - 1), & W_{2m+1}^{(2)} &\equiv -(2m - 1)k + 2m && \pmod{(4k(k - 1))} \\
W_{2m}^{(3)} &\equiv (2m + 1)k - 2m, & W_{2m+1}^{(3)} &\equiv -2mk + 2m + 1 && \pmod{(4k(k - 1))} \\
W_{2m}^{(4)} &\equiv (1 - 2m)k + 2m, & W_{2m+1}^{(4)} &\equiv 2mk - (2m + 1) && \pmod{(4k(k - 1))} \\
W_{2m}^{(5)} &\equiv (2m + 2)k - (2m + 1), & W_{2m+1}^{(5)} &\equiv (-2m - 1)k + 2m && \pmod{(4k(k - 1))} \\
W_{2m}^{(6)} &\equiv (2 - 2m)k + (2m - 1), & W_{2m+1}^{(6)} &\equiv (2m + 1)k - 2m && \pmod{(4k(k - 1))}
\end{aligned}$$

Proof. All the claims are proven inductively. We first prove that the sequence $(|W_m^{(1)}|)_m$ is increasing.

For $m = 1$, the inequality $|W_1| \geq |W_0|$ holds since $|4k^2 - k - 2| \geq 4|k|^2 - |k| - 2 \geq 1 = |W_0|$ for $|k| \geq 1$. From $W_{m+1} = 2(4k^2 - 2k - 1)W_m - W_{m-1}$, it follows that $|W_{m+1}| \geq (8|k|^2 - 4|k| - 2)|W_m| - |W_{m-1}| \geq (8|k|^2 - 4|k| - 3)|W_m| + |W_m| - |W_{m-1}| \geq (8|k|^2 - 4|k| - 3)|W_m|$.

This directly shows not only that the sequence of absolute values is increasing, but also that $|W_m| \geq (8|k|^2 - 4|k| - 3)^{m-1}$ for all $m \geq 0$. Likewise, this holds for $W_m^{(i)}$ for all $i = 1, \dots, 6$.

Inductively one shows that $W_{2m}^{(1)} \equiv -2mk + 2m + 1 \pmod{4k(k - 1)}$ and $W_{2m+1}^{(1)} \equiv (2m + 3)k - 2m - 2 \pmod{4k(k - 1)}$. The inductive basis, for $m = 0$, was computed before the lemma statement. If we assume that $W_{2m}^{(1)} \equiv -2mk + 2m + 1 \pmod{4k(k - 1)}$ and $W_{2m+1}^{(1)} \equiv (2m + 3)k - 2m - 2 \pmod{4k(k - 1)}$, then

$$\begin{aligned}
W_{2m+2}^{(1)} &= 2(4k^2 - 2k - 1)W_{2m+1} - W_{2m} \\
&\equiv 2(2k - 1) \cdot ((2m + 3)k - 2m - 2) - (-2mk + 2m + 1) \\
&= (2m + 3)(4k^2 - 2k) - (2m + 2)(4k - 2) + 2mk - 2m - 1 \\
&\equiv (2m + 3) \cdot 2k - 8mk + 4m - 8k + 4 + 2mk - 2m - 1 \\
&= -2(m + 1)k + 2m + 3 \pmod{4k(k - 1)}.
\end{aligned}$$

Similarly, by using this claim, one gets that

$$\begin{aligned}
W_{2m+3}^{(1)} &= 2(4k^2 - 2k - 1)W_{2m+2} - W_{2m+1} \\
&\equiv 2(2k - 1)(-2(m + 1)k + 2m + 3) - ((2m + 3)k - 2m - 2) \\
&= -2(m + 1)(4k^2 - 2k) + 2(2k - 1)(2m + 3) - (2m + 3)k + 2m + 2 \\
&\equiv -2(m + 1) \cdot 2k + 8mk + 12k - 4m - 6 - 2mk - 3k + 2m + 2 \\
&= (2m + 5)k - 2m - 4 \pmod{4k(k - 1)}.
\end{aligned}$$

This shows the congruence claims for $(W_m^{(1)})_m$, while the claims stated for the other sequences are proven completely analogously. \square

Now we observe the even indices case: $W_{2m}^{(1)} = V_{2n} \Rightarrow -2mk + 2m + 1 \equiv 1 \pmod{4k(k - 1)}$ so $4k^2 - 4k$ divides $2mk - 2m$, i. e. $2k(k - 1) | m(k - 1)$ and, most importantly, $2k \mid m$. Analogously,

$W_{2m} = V_{2m+1} \equiv 2k - 1 \pmod{4k(k-1)}$ implies that $2k \mid m + 1$. Any of these conclusions, $2k \mid m$ and $2k \mid m + 1$, implies that

$$(5.1) \quad m \geq 2|k| - 1,$$

unless m is such that the corresponding multiple of $2k$ is actually 0. Therefore, if $m \neq 0, -1$, then

$$|x| \geq (8|k|^2 - 4|k| - 3)^{4|k|-3}.$$

For odd indices in the sequence $(W_m^{(1)})$, $(2m+3)k - 2m - 2 \equiv 1 \pmod{4k(k-1)}$ implies that $4k(k-1) \mid (2m+3)(k-1)$ and $4k \mid 2m+3$, which is obviously impossible. Analogously, $(2m+3)k - 2m - 2 \equiv 1 \pmod{4k(k-1)}$ implies $4k \mid 2m+1$, a contradiction.

In the same way, one gets similar conclusions for the remaining sequences $(W_m^{(i)})_m$ ($i = 2, \dots, 6$): for one case (even/odd index) there is a contradiction, while the other case implies that $2k \mid m$ or $2k \mid m \pm 1$. In any case, $m \geq 2|k| - 1$ if $m \notin \{-1, 0, 1\}$ and the same lower bound holds. We have proven the following result.

Proposition 5.2. *If (x, y, z) is the solution of the system*

$$(4.1) \quad (k+1)x^2 - (k-1)y^2 = 2,$$

$$(4.2) \quad (16k^3 - 4k)x^2 - (k-1)z^2 = 16k^3 - 5k + 1,$$

for $|k| > 17$ and $x \notin \{1, k, 2k-1, 8k^3 - 4k^2 - 4k + 1\}$, then $|x| \geq (8|k|^2 - 4|k| - 3)^{4|k|-3}$.

We note here that the exceptions $x = 1$, $x = k$, $x = 2k - 1$ and $x = 8k^3 - 4k^2 - 4k + 1$ correspond to the indices $m = 0$ and $m = 1$, i. e. when $2k \mid m$ does not imply that $m \geq 2|k|$.

6. THE PROBLEM OF APPLYING JADRIJEVIĆ–ZIEGLER THEOREM

There are two essentially different systems we can attempt to solve in this problem. One is given in Proposition 5.2 and Jadrijević–Ziegler theorem [12] cannot be applied here because its conditions are not satisfied. The second system has coefficient $16k^3 - 4k$ on left-hand side of both of the equations – we will show that, while the conditions are satisfied, this theorem cannot give us a useful result.

First, we focus on the system already given.

Lemma 6.1. *If (x, y, z) is a solution of the system of equations (4.1) and (4.2), and $\theta_1^{(1)} = \pm \sqrt{\frac{k+1}{k-1}}$,*

$\theta_1^{(2)} = -\theta_1^{(1)}$, $\theta_2^{(1)} = \pm \sqrt{\frac{4k^2 - 1}{4k(k-1)}}$, $\theta_2^{(2)} = -\theta_2^{(1)}$, where signs are chosen in such a way that

$$\left| \theta_1^{(1)} - \frac{y}{x} \right| \leq \left| \theta_1^{(2)} - \frac{y}{x} \right| \quad \text{and} \quad \left| \theta_2^{(1)} - \frac{z}{4kx} \right| \leq \left| \theta_2^{(2)} - \frac{z}{4kx} \right|,$$

then

$$\left| \theta_1^{(1)} - \frac{4ky}{4kx} \right| \leq \frac{2}{\sqrt{|k^2 - 1|}} \cdot \frac{1}{|x|^2}, \quad \text{and}$$

$$\left| \theta_2^{(1)} - \frac{z}{4kx} \right| \leq \frac{|16k^3 - 5k + 1|}{8\sqrt{|4k^6 - 4k^5 - k^4 + k^3|}} \cdot \frac{1}{|x|^2}.$$

Proof. The first inequality, $\left| \theta_1^{(1)} - \frac{4ky}{4kx} \right| \leq \frac{2}{\sqrt{|k^2 - 1|}} \cdot \frac{1}{|x|^2}$, was already obtained in [9]. In the same manner,

$$\begin{aligned} \left| \theta_2^{(1)} - \frac{z}{4kx} \right| &= \left| (\theta_2^{(1)})^2 - \frac{z^2}{16k^2x^2} \right| \cdot \left| \theta_2^{(1)} + \frac{z}{4kx} \right|^{-1} \\ &= \left| \frac{1}{16k^2} \left(\frac{16k^3 - 4k}{k - 1} - \frac{z^2}{x^2} \right) \right| \cdot \left| \theta_2^{(2)} - \frac{z}{4kx} \right|^{-1} \\ (6.1) \quad &\stackrel{(4.2)}{=} \frac{|16k^3 - 5k + 1|}{|16k^3 - 16k^2|} \cdot \frac{1}{|x|^2} \cdot \left| \theta_2^{(2)} - \frac{z}{4kx} \right|^{-1} \end{aligned}$$

Furthermore, because of the way the signs were chosen,

$$\left| \theta_2^{(2)} - \frac{z}{4kx} \right| \geq \frac{1}{2} \left(\left| \theta_2^{(1)} - \frac{z}{4kx} \right| + \left| \theta_2^{(2)} - \frac{z}{4kx} \right| \right) \geq \frac{1}{2} |\theta_2^{(1)} - \theta_2^{(2)}| = \left| \sqrt{\frac{4k^2 - 1}{4k^2 - 4k}} \right|$$

Plugging in (6.1), we get

$$\left| \theta_2^{(1)} - \frac{z}{4kx} \right| \leq \frac{|16k^3 - 5k + 1|}{|16k^3 - 16k^2|} \cdot \left| \sqrt{\frac{4k(k-1)}{4k^2 - 1}} \right| \cdot \frac{1}{|x|^2} = \frac{|16k^3 - 5k + 1|}{8\sqrt{|4k^6 - 4k^5 - k^4 + k^3|}} \cdot \frac{1}{|x|^2}.$$

□

Now we want to apply the following theorem [12].

Theorem 6.2 ([12, Theorem 7.1]). *Let $\theta_i = \sqrt{1 + \frac{a_i}{T}}$, $i = 1, 2$ where $a_1 \neq a_2$ and T are in the ring of integers of an imaginary quadratic field K . Let $|T| > M = \max\{|a_1|, |a_2|\}$,*

$$L = \frac{27}{16|a_1|^2|a_2|^2|a_1 - a_2|^2} (|T| - M)^2 > 1.$$

Then

$$\max \left\{ \left| \theta_1 - \frac{p_1}{q} \right|, \left| \theta_2 - \frac{p_2}{q} \right| \right\} > c|q|^{-\lambda},$$

holds for all algebraic integers $p_1, p_2, q \in K$, where $\lambda = 1 + \frac{\log P}{\log L}$, $c^{-1} = 4pP(\max\{1, 2l\})^{\lambda-1}$,

$$l = \frac{27}{64} \frac{|T|}{|T| - M}, p = \sqrt{\frac{2|T| + 3M}{2|T| - 2M}}, P = 16 \frac{|a_1|^2 |a_2|^2 |a_1 - a_2|^2}{\min\{|a_1|, |a_2|, |a_1 - a_2|\}^3} (2|T| + 3M).$$

Writing $\theta_1 = \sqrt{\frac{k+1}{k-1}} = \sqrt{1 + \frac{2}{k-1}}$, $\theta_2 = \sqrt{\frac{4k^2-1}{4k(k-1)}} = \sqrt{1 + \frac{4k-1}{4k^2-4k}}$, we see that, to get

the same denominators, we need to write θ_1 as $\theta_1 = \sqrt{1 + \frac{8k}{4k^2-4k}}$. Hence $a_1 = 8k, a_2 = 4k-1, T = 4k^2 - 4k$. Then $M = \max\{|a_1|, |a_2|\} = 8|k|$. The inequality $|k-1| \geq |k-1| > 2$ holds for $|k| > 3$, so $|T| = 4|k||k-1| > 8|k| = M$.

Unfortunately, $L = \frac{27}{16 \cdot (8|k|)^2 \cdot |4k-1|^2 \cdot |4k+1|^2} (4|k^2-k| - 8|k|)^2$ is, for large k , less than 1 (since the degree of $|k|$ is 6 in the denominator and 4 in the numerator), while the condition of the theorem is $L > 1$. Therefore, we cannot directly apply this theorem.

On the other hand, we can attempt to solve the following system

$$(6.2) \quad (16k^3 - 4k)y^2 - (k+1)z^2 = 16k^3 - 5k - 1,$$

$$(4.2) \quad (16k^3 - 4k)x^2 - (k-1)z^2 = 16k^3 - 5k + 1$$

and define ϑ_1, ϑ_2 as

$$\vartheta_1^2 = 1 + \frac{1}{(k-1)(16k^3-4k)}, \quad \vartheta_2^2 = 1 + \frac{1}{(k+1)(16k^3-4k)},$$

where the signs of ϑ_1 and ϑ_2 are chosen in the same manner as in Lemma 6.1. In that case, by the notation of Jadrijević-Ziegler theorem,

$$a_1 = k+1, \quad a_2 = k-1, \quad T = (k^2-1)(16k^3-4k).$$

Remark 7.2 from [12] shows that the condition $L > 1$ is fulfilled whenever $|T| > (4M)^3$. Here, this inequality $|(k^2-1)(16k^3-4k)| > |4(k+1)|^3$ holds for $k \geq 3.21$.

Now we need to show that ϑ_1 and ϑ_2 can be approximated by the quotient of solutions (up to multiplication by some element of $\mathbb{Q}[i]$). More precisely, we will bound $\left| \vartheta_1 - \frac{sx}{(k-1)z} \right|$ in the following lemma, where $s^2 = (4k^2 - 2k - 1)^2$, and a similar expression for ϑ_2 .

Lemma 6.3. For $|k| \geq 5$,

$$\max \left\{ \left| \vartheta_1^{(1)} - \frac{s(k+1)x}{(k-1)(k+1)z} \right|, \left| \vartheta_2^{(1)} - \frac{t(k-1)y}{(k-1)(k+1)z} \right| \right\} < 40|k|^2|z|^{-2},$$

where $t = 4k^2 + 2k - 1$.

Proof.

$$\begin{aligned}
& \left| \vartheta_1^{(1)} - \frac{sx}{(k-1)z} \right| = \\
& = \left| (\vartheta_1^{(1)})^2 - \frac{s^2x^2}{(k-1)^2z^2} \right| \cdot \left| \vartheta_1^{(1)} + \frac{sx}{(k-1)z} \right|^{-1} \\
& = \left| 1 + \frac{1}{(k-1)(16k^3-4k)} - \frac{s^2x^2}{(k-1)^2z^2} \right| \cdot \left| \vartheta_1^{(2)} - \frac{sx}{(k-1)z} \right|^{-1} \\
& = \left| \frac{(k-1)^2(16k^3-4k)z^2 + (k-1)z^2 - ((k-1)(16k^3-4k)+1)(16k^3-4k)x^2}{(k-1)^2(16k^3-4k)z^2} \right| \\
& \quad \cdot \left| \vartheta_1^{(2)} - \frac{sx}{(k-1)z} \right|^{-1} \\
& = \left| \frac{(k-1)(16k^3-4k)((k-1)z^2 - (16k^3-4k)x^2) + (k-1)z^2 - (16k^3-4k)x^2}{(k-1)^2(16k^3-4k)z^2} \right| \\
& \quad \cdot \left| \vartheta_1^{(2)} - \frac{sx}{(k-1)z} \right|^{-1} \\
& = \left| \frac{((k-1)(16k^3-4k)+1)((k-1)z^2 - (16k^3-4k)x^2)}{(k-1)^2 - (16k^3-4k)z^2} \right| \cdot \left| \vartheta_1^{(2)} - \frac{sx}{(k-1)z} \right|^{-1} \\
& \stackrel{(4.2)}{=} \left| \frac{s^2((k-1) - (16k^3-4k))}{(k-1)^2(16k^3-4k)} \right| \cdot \left| \vartheta_1^{(2)} - \frac{sx}{(k-1)z} \right|^{-1} \cdot |z|^{-2}.
\end{aligned}$$

Since $t^2 = (k+1)(16k^3-4k)+1 = (4k^2+2k-1)^2$, in the same way it follows that

$$\left| \vartheta_2^{(1)} - \frac{ty}{(k+1)z} \right| = \left| \frac{t^2((k+1) - (16k^3-4k))}{(k+1)^2(16k^3-4k)} \right| \cdot \left| \vartheta_2^{(2)} - \frac{ty}{(k+1)z} \right|^{-1} \cdot |z|^{-2}.$$

Analogously as before,

$$\begin{aligned}
\left| \vartheta_1^{(2)} - \frac{sx}{(k-1)z} \right| & \geq \frac{1}{2} |\vartheta_1^{(1)} - \vartheta_1^{(2)}| = \left| \sqrt{1 + \frac{1}{(k-1)(16k^3-4k)}} \right| = \left| \frac{4k^2-2k-1}{\sqrt{(k-1)(16k^3-4k)}} \right|, \\
\left| \vartheta_2^{(2)} - \frac{ty}{(k+1)z} \right| & \geq \frac{1}{2} |\vartheta_2^{(1)} - \vartheta_2^{(2)}| = \left| \sqrt{1 + \frac{1}{(k+1)(16k^3-4k)}} \right| = \left| \frac{4k^2+2k-1}{\sqrt{(k+1)(16k^3-4k)}} \right|.
\end{aligned}$$

Now

$$\begin{aligned}
& \left| \frac{s^2((k-1) - (16k^3 - 4k))}{(k-1)^2(16k^3 - 4k)} \right| \cdot \left| \vartheta_1^{(2)} - \frac{sx}{(k-1)z} \right|^{-1} \leq \\
& \left| \frac{s^2((k-1) - (16k^3 - 4k))}{(k-1)^2(16k^3 - 4k)} \right| \cdot \left| \frac{4k^2 - 2k - 1}{\sqrt{(k-1)(16k^3 - 4k)}} \right|^{-1} \\
& = \left| \frac{(4k^2 - 2k - 1)^2(16k^3 - 5k + 1)}{(k-1)^2(16k^3 - 4k)} \right| \cdot \left| \frac{\sqrt{(k-1)(16k^3 - 4k)}}{4k^2 - 2k - 1} \right| \\
& = \left| \frac{(4k^2 - 2k - 1)(16k^3 - 5k + 1)}{(k-1)^2(16k^3 - 4k)} \right| \cdot |\sqrt{(k-1)(16k^3 - 4k)}| \\
& = \left| \frac{64k^5 - 32k^4 - 36k^3 + 14k^2 + 3k - 1}{16k^5 - 32k^4 + 12k^3 + 8k^2 - 4k} \right| \cdot \sqrt{|k-1| \cdot |16k^3 - 4k|} \\
& \leq \frac{64|k|^5 + 32|k|^4 + 36|k|^3 + 14|k|^2 + 3|k| + 1}{16|k|^5 - 32|k|^4 - 12|k|^3 - 8|k|^2 - 4|k|} \sqrt{16|k|^4 + 16|k|^3 + 4|k|^2 + |k|} \\
& \leq \frac{64|k|^5 + 32|k|^4 + 36|k|^3 + 14|k|^2 + 3|k| + 1}{16|k|^5 - 32|k|^4 - 12|k|^3 - 8|k|^2 - 4|k|} \cdot (4|k|^2 + 2|k| + 1)
\end{aligned}$$

The last used inequality is easily proven by squaring it. It suffices to show that $(64|k|^5 + 32|k|^4 + 36|k|^3 + 14|k|^2 + 3|k| + 1)(4|k|^2 + 2|k| + 1) \leq 40|k|^2(16|k|^5 - 32|k|^4 - 12|k|^3 - 8|k|^2 - 4|k|)$, which is equivalent to $384|k|^7 - 1536|k|^6 - 752|k|^5 - 480|k|^4 - 236|k|^3 - 24|k|^2 - 5|k| - 1 \geq 0$. Since $384|k|^7 \geq 1920|k|^6$ for $|k| \geq 5$, so it suffices to show that $384|k|^6 - 752|k|^5 - 480|k|^4 - 236|k|^3 - 24|k|^2 - 5|k| - 1 \geq 0$. By repeating this argument, we get the proof of the desired inequality. \square

We now show that $2l = 2 \cdot \frac{27}{64} \frac{|T|}{|T| - M} < 1$. This is equivalent to $27|T| < 32|T| - 32M$, i. e. $32M < 5|T|$. Since M is the larger among the numbers $|k-1|$ and $|k+1|$, and both of them are less or equal to $|k|+1$ (by the triangle inequality), it follows that $M \leq |k|+1$. Therefore, we can show that $32(|k|+1) < 5|16k^5 - 20k^3 + 4|$, which holds for $|k| \geq 1.33$. Namely, $5|16k^5 - 20k^3 + 4| \geq 80|k|^5 - 100|k|^3 - 20$, so it suffices to show that $80|k|^5 - 100|k|^3 - 32|k| - 52 > 0$, which holds for k with large absolute value.

$$\text{Now } c = \frac{1}{4pP}, L = \frac{27(|T|-M)^2}{64|k^2-1|^2}, p = \sqrt{1 + \frac{5M}{2|T|-2M}}, P = 8(2|T| + 3M)|k^2 - 1|^2, q = (k-1)(k+1)z.$$

If we try to apply the Jadrijević-Ziegler theorem, then

$$\lambda = 1 + \frac{\log 8 + \log(2|T| + 3M) + 2 \log |k^2 - 1|}{\log 27 + 2 \log (|T| - M) - \log 64|k^2 - 1|^2},$$

and let

$$\text{Max} = \max \left\{ \left| \vartheta_1^{(1)} - \frac{s(k+1)x}{(k-1)(k+1)z} \right|, \left| \vartheta_2^{(1)} - \frac{t(k-1)y}{(k-1)(k+1)z} \right| \right\}.$$

Then

$$\text{Max} > \frac{1}{4pP}|q|^{-\lambda} = \frac{\sqrt{2|T| - 2M}}{32 \sqrt{2|T| + 3M}(2|T| + 3M)|k^2 - 1|^2}|q|^{-\lambda}.$$

Since $M < \frac{5}{32}|T|$, it follows that $2|T| - 2M > \frac{27}{16}|T|$, and $2|T| + 3M < \frac{79}{32}|T| < \frac{5}{2}|T|$. Hence

$$\text{Max} > \frac{\sqrt{\frac{27}{16}|T|}}{32 \cdot \frac{5}{2} \sqrt{\frac{5}{2}|T|^{\frac{3}{2}}|k^2 - 1|^2}}|q|^{-\lambda} = \frac{3\sqrt{3}}{160\sqrt{10}} \frac{|T|^{-1}|q|^{-\lambda}}{|k^2 - 1|^2} = C|k^2 - 1|^{-\lambda-3}|16k^3 - 4k|^{-1}|z|^{-\lambda},$$

where $C = \frac{3\sqrt{3}}{160\sqrt{10}}$. Now we can conclude that $C|k^2 - 1|^{-\lambda-3}|16k^3 - 4k|^{-1}|z|^{-\lambda} < 40|k|^2|z|^{-2}$, i. e.

$$|z|^{2-\lambda} < \frac{40}{C}|k|^2|k^2 - 1|^{\lambda+3}|16k^3 - 4k| = \frac{6400\sqrt{10}}{3\sqrt{3}}|k|^2|k^2 - 1|^{\lambda+3}|16k^3 - 4k|.$$

This inequality can be used to bound the magnitude of solution $|z|$ when $\lambda < 2$, because the left-hand side is then a positive power of $|z|$. The proof for lower bound on $|x|$ is easily modified for $|z|$. It is not hard to see that $|z| \geq |x|$, so we could use the same lower bound. Since this lower bound is exponential in $|k|$, if λ were less than 2, then we would get a polynomial upper bound for $|z|$ and juxtaposition of these two bounds would give us the upper bound for $|k|$. Unfortunately, $\lambda > 2$ here. Namely, this claim is equivalent to $P > L$ and $8(2|T| + 3M)|k^2 - 1|^2 > \frac{27(|T| - M)^2}{64|k^2 - 1|^2}$, and $512(2|(k^2 - 1)(16k^3 - 4k)| + 3M)|k^2 - 1|^4 > 27(|(k^2 - 1)(16k^3 - 4k)| - M)^2$. Since $M = \max\{|k - 1|, |k + 1|\}$ is linear in k , we can already see that the degree of k is greater in the left-hand side (13 > 10). More precisely, left-hand side is $512(2|(k^2 - 1)(16k^3 - 4k)| + 3M)|k^2 - 1|^4 \geq 512(32|k|^5 - 40|k|^3 - 11|k| - 3)(|k^2 - 1|^4)$, while $27(|(k^2 - 1)(16k^3 - 4k)| - M)^2 < 27(16|k|^5 + 20|k|^3 + 4|k|)^2$. It suffices to check that

$$\begin{aligned} &16384|k|^{13} - 86016|k|^{11} - 6912|k|^{10} + 174592|k|^9 - 18816|k|^8 - 165888|k|^7 - 8112|k|^6 + \\ &+ 64512|k|^5 - 13536|k|^4 + 2048|k|^3 + 5712|k|^2 - 5632|k| - 1536 > 0, \end{aligned}$$

which holds for $|k| \geq 1.82$.

To conclude, the gap between $16k^3 - 4k$ and $k + 1$ is not large enough for exponent λ to be less than 2, and this makes it unlikely to use the usual approach by Diophantine approximation.

We note here that the similar problem of extending $D(4)$ -triple $\{k' - 2, k' + 2, 4(k')^3 - 4k'\}$ in rational integers was studied in [2]. For even $k' = 2k$, dividing by 2, we get $D(1)$ -triples having the same form as the triples studied in this paper. In [2], problem was solved using a similar method we tried to apply here. An amelioration of the analogous theorem in \mathbb{Z} was proven there for a specific situation (where numerators under the square root in θ_i equal exactly $k - 2$ and $k + 2$, while the denominator is divisible by $k^2 - 4$).

7. APPLICATION OF LINEAR FORMS IN LOGARITHMS TO THE FAMILY $\{k - 1, k + 1, 16k^3 - 4k\}$

We continue dealing with the extensibility problem of Diophantine triples $\{k - 1, k + 1, 16k^3 - 4k\}$. Sequence $(V_n)_n$ is defined as in (4.3), while $(W_m^{(i)})_m$ is defined in Lemma 4.3. Let us remind ourselves that for this family, $a = k - 1, b = k + 1, c = 16k^3 - 4k$ and $r = k, s = 4k^2 - 2k - 1, t = 4k^2 + 2k + 1$.

Lemma 7.1. *If $V_n = \pm W_m^{(j)}$ for some $j, m, n \in \mathbb{N}_0$ and $|k| > 2.5$, then $m \leq n \leq 3m + 2$.*

Proof. Recurrence relations and Lemma 5.1 inductively imply the following inequalities

$$(2|k| - 1)^n \leq |V_n| \leq (2|k| + 1)^n$$

$$(8|k|^2 - 4|k| - 3)^{m-1} \leq |W_m^{(j)}| \leq (8|k|^2 + 4|k| + 3)^{m+1}.$$

If $V_n = W_m$, then $(2|k| + 1)^n \geq (8|k|^2 - 4|k| - 3)^{m-1}$, so $n \geq m$. If we assume the contrary, $n \leq m - 1$, then $8|k|^2 - 4|k| - 3 \leq 2|k| + 1$, which creates a contradiction when $|k| > 2.5$.

We now assume $n \geq 3m + 3$. From $V_n = W_m$ it follows that $(8|k|^2 + 4|k| + 3)^{m+1} \geq (2|k| - 1)^n \geq (2|k| - 1)^{3m+3}$, so $8|k|^2 + 4|k| + 3 > (2|k| - 1)^3 = 8|k|^3 - 12|k|^2 + 6|k| - 1$. This implies that $-2(4|k|^3 - 10|k|^2 + |k| - 2) > 0$, which is impossible for $|k| > 2.5$. \square

By solving the recurrence relations defining (V_n) and (W_m) , we get that

$$V_n = \frac{\sqrt{k+1} + \sqrt{k-1}}{2\sqrt{k+1}}(k + \sqrt{k^2 - 1})^n + \frac{\sqrt{k+1} - \sqrt{k-1}}{2\sqrt{k+1}}(k - \sqrt{k^2 - 1})^n,$$

$$W_m = \frac{1}{2\sqrt{16k^3 - 4k}}((x_1 \sqrt{16k^3 - 4k} + z_1 \sqrt{k-1})(4k^2 - 2k - 1 + \sqrt{(16k^3 - 4k)(k-1)})^m +$$

$$+ (x_1 \sqrt{16k^3 - 4k} - z_1 \sqrt{k-1})(4k^2 - 2k - 1 - \sqrt{(16k^3 - 4k)(k-1)})^m).$$

Let $P' = \frac{1}{\sqrt{c}}(x_1 \sqrt{c} + z_1 \sqrt{a})(s + \sqrt{ac})^m$ i $Q' = \frac{1}{\sqrt{b}}(\sqrt{a} + \sqrt{b})(r + \sqrt{ab})^n$. We remark that $Q' \neq Q$ and $P' = \sqrt{\frac{a}{c}}P$. However, with $m \geq 3$, we have the same bounds on Q' and $|P'| - |Q'|$. They are obtained in a similar manner:

$$|Q'| = \frac{1}{\sqrt{|b|}}|\sqrt{a} + \sqrt{b}| \cdot |r + \sqrt{ab}|^n \geq \frac{|b-a|}{\sqrt{|b|}} \cdot \frac{1}{|\sqrt{b} - \sqrt{a}|} |\sqrt{ab}|^3$$

$$\geq \frac{2}{\sqrt{|b|}} \cdot \frac{1}{|\sqrt{|b|} + |\sqrt{a}|} |ab|^{3/2} \geq 12 \frac{|b|}{|a|},$$

since $|a|^{5/2} \geq 6(\sqrt{|b|} + \sqrt{|a|})$, i. e. $|k+1|^{5/2} \geq 6(\sqrt{|k+1|} + \sqrt{|k-1|})$, which holds for $|k| \geq 4.846$. If $|k| \geq 23$, this implies that $|Q'| \geq 11$ since $12 \frac{|k+1|}{|k-1|} \geq 12 \frac{|k|-1}{|k|+1} \geq 11$, which is equivalent to $|k| \geq 23$.

Similarly, it holds that $|P'| \geq 12$ so

$$\begin{aligned} \left| |P'| - |Q'| \right| &\leq \left| \frac{c-a}{c}(P')^{-1} - \frac{b-a}{b}(Q')^{-1} \right| \leq \left| 1 - \frac{a}{c} \right| \frac{1}{|P'|} + \left| 1 - \frac{a}{b} \right| \frac{1}{|Q'|} \\ &\leq \frac{1}{12} \left| 1 - \frac{a}{c} \right| + \frac{1}{11} \frac{2}{|b|} < \frac{5}{48} + \frac{5}{48} = \frac{5}{24}, \end{aligned}$$

for $|b| \geq |k| - 1 > \frac{96}{55}$. Therefore, the conclusion of Lemma 2.6 holds for the linear form $\Gamma = \log \Lambda' = \log \frac{|P'|}{|Q'|}$ as well.

7.1. Minimal polynomials. Let $k = \mu + iv$ and

$$\begin{aligned} \alpha_1 &= |k + \sqrt{k^2 - 1}|, \\ \alpha_2 &= |4k^2 - 2k - 1 + \sqrt{(16k^3 - 4k)(k - 1)}| i \\ \alpha_3 &= \left| \frac{\sqrt{16k^3 - 4k}(\sqrt{k - 1} + \sqrt{k + 1})}{\sqrt{k + 1}(x_1 \sqrt{16k^3 - 4k} + z_1 \sqrt{k + 1})} \right|. \end{aligned}$$

The minimal polynomial for α_1 is $p_1(x) = x^8 - 4(\mu^2 + \nu^2)x^6 + (8\mu^2 - 8\nu^2 - 2)x^4 - 4(\mu^2 + \nu^2)x^2 + 1$, according to [9]. In the same paper, it was shown that $h(\alpha_1) \leq \frac{1}{4} \log(2|k| + 1)$.

The minimal polynomial for α_2 is determined with the help of Mathematica [14],

$$\begin{aligned} p_2(x) &= x^8 - 4 \left((16(\mu^2 + \nu^2) - 16\mu - 4)(\mu^2 + \nu^2) + 16\nu^2 + 4\mu + 1 \right) x^6 + \\ &\quad + \left((128\mu^4 - 128\mu^3 - 32\mu^2 + 32\mu + 6) + (-768\mu^2 + 384\mu + 32)\nu^2 + 128\nu^4 \right) x^4 \\ &\quad - 4 \left((16(\mu^2 + \nu^2) - 16\mu - 4)(\mu^2 + \nu^2) + 16\nu^2 + 4\nu + 1 \right) x^2 + 1. \end{aligned}$$

Polynomial $p_2(x)$ has the following zeroes:

$$\begin{aligned} x_{1,2} &= \pm \alpha_2, & x_{5,6} &= \pm \sqrt{|s|^2 - |ac| + \sqrt{(|s|^2 - |ac|)^2 - 1}}, \\ x_{3,4} &= \pm |s - \sqrt{ac}|, & x_{7,8} &= \pm \sqrt{|s|^2 - |ac| - \sqrt{(|s|^2 - |ac|)^2 - 1}}, \end{aligned}$$

and $|x_i| = 1$ for $i = 5, 6, 7, 8$. It follows that

$$h(\alpha_2) \leq \frac{1}{8} \log |x_1||x_2| = \frac{1}{4} \log |4k^2 - 2k - 1 + \sqrt{(16k^3 - 4k)(k - 1)}|,$$

which implies $h(\alpha_2) \leq \frac{1}{4} \log |9k^2| = \frac{1}{2} \log 3|k|$. Polynomial $p_2(x)$ has the following zeroes $x_{1,2} = \pm \alpha_2$, $x_{3,4} = \pm |s - \sqrt{ac}|$, $x_{5,6} = \pm \sqrt{|s|^2 - |ac| + \sqrt{(|s|^2 - |ac|)^2 - 1}}$ and $x_{7,8} = \pm \sqrt{|s|^2 - |ac| - \sqrt{(|s|^2 - |ac|)^2 - 1}}$, and $|x_i| = 1$ for $i = 5, 6, 7, 8$. This implies that

$$h(\alpha_2) \leq \frac{1}{8} \log |x_1||x_2| = \frac{1}{4} \log |4k^2 - 2k - 1 + \sqrt{(16k^3 - 4k)(k - 1)}|,$$

and, consequently $h(\alpha_2) \leq \frac{1}{4} \log |9k^2| = \frac{1}{2} \log 3|k|$.

7.2. Bounding the conjugates of α_3 .

Lemma 7.2. *If $|k| \geq 10^7$, then, for all conjugates α'_3 of α_3 , it holds that $|\alpha'_3| \leq |k|^4$.*

Proof. One can guess the minimal polynomial for α_3 and all conjugates. The first eight are $x'_{1,2} = \pm\alpha_3$,

$$x_{3,4} = \pm \left| \frac{\sqrt{c}(\sqrt{b} + \sqrt{a})}{\sqrt{b}(x_1 \sqrt{c} - z_1 \sqrt{a})} \right|, x_{5,6} = \pm \left| \frac{\sqrt{c}(\sqrt{b} - \sqrt{a})}{\sqrt{b}(x_1 \sqrt{c} - z_1 \sqrt{a})} \right|, x_{7,8} = \pm \left| \frac{\sqrt{c}(\sqrt{b} - \sqrt{a})}{\sqrt{b}(x_1 \sqrt{c} - z_1 \sqrt{a})} \right|.$$

Furthermore, x_9, \dots, x_{12} are zeroes of

$$q_1(x) = x^4 - 2 \left| \frac{c}{b(x_1 \sqrt{c} + z_1 \sqrt{a})^2} \right| (|b| - |a|)x^2 + \left| \frac{c(b-a)}{b(x_1 \sqrt{c} + z_1 \sqrt{a})^2} \right|^2,$$

x_{13}, \dots, x_{16} of

$$q_2(x) = x^4 - 2 \left| \frac{c}{b(x_1 \sqrt{c} - z_1 \sqrt{a})^2} \right| (|b| - |a|)x^2 + \left| \frac{c(b-a)}{b(x_1 \sqrt{c} - z_1 \sqrt{a})^2} \right|^2,$$

the next eight of

$$q_3(x) = x^4 - 2 \left| \frac{c(\sqrt{b} + \sqrt{a})^2}{b(c-a)^2} \right| (|cx_1^2| - |az_1^2|)x^2 + \left| \frac{c(\sqrt{b} - \sqrt{a})^2}{b(c-a)} \right|^2 \quad \text{and}$$

$$q_4(x) = x^4 - 2 \left| \frac{c(\sqrt{b} - \sqrt{a})^2}{b(c-a)^2} \right| (|cx_1^2| - |az_1^2|)x^2 + \left| \frac{c(\sqrt{b} - \sqrt{a})^2}{b(c-a)} \right|^2,$$

then

$$q_5(x) = x^4 - 2 \left| \frac{c}{b(c-a)^2} \right| (|x_1 \sqrt{bc} + z_1 a|^2 - |x_1 \sqrt{ac} + z_1 \sqrt{ab}|^2)x^2 + \left| \frac{c(b-a)}{b(c-a)} \right|^2$$

and finally

$$q_6(x) = x^4 - 2 \left| \frac{c}{b(c-a)^2} \right| (|x_1 \sqrt{bc} - z_1 a|^2 - |x_1 \sqrt{ac} - z_1 \sqrt{ab}|^2)x^2 + \left| \frac{c(b-a)}{b(c-a)} \right|^2.$$

This suffices to find the bound we need here. Namely, the zero of the monic polynomial is bounded from above by the sum of the absolute values of its coefficients. For this polynomial, we can see that the coefficients have at most the order of $|c|^2 \cdot |a|$ (or $\cdot |b|$). More precisely, we will show that all the coefficients of x^2 -terms are less than $3|k|^7$, while all the free coefficients are less than $1025|k|^7$ for k large enough.

The coefficient of x^2 in q_1 and q_2 is less than or equal to $2|c|(|b| + |a|) \leq 2|16k^3 - 4k|(2|k| + 2) \leq |k|^5$ for $|k| \geq 65$. This type of claim is proven as earlier in the paper, by using the triangle inequality and analysing the obtained functions of $|k|$. The coefficient of x^2 in q_3 and q_4 is less than or equal to

$$\frac{2|c|(\sqrt{|a|} + \sqrt{|b|})(|cx_1^2| + |az_1^2|)}{|b|^2|c-a|} \leq \frac{2|16k^3 - 4k| \cdot 2\sqrt{|k|+1}(150|k|^5 + 65)}{(|k|-1)^2(16|k|^3 - 5|k|-1)} < |k|^4 \quad \text{for } |k| \geq 1.45 \cdot 10^6.$$

Similarly, the coefficient of x^2 in q_5 and q_6 is less than $3|k|^7$ for $|k| \geq 2$.

The free coefficients are less than $|16k^3 - 4k|^2(2\sqrt{|k|+1})^2 \leq 1025|k|^7$ for $|k| \geq 1025$.

Therefore, $|\alpha'_3|^2 \leq 1028|k|^7$ for every conjugate α'_3 , i. e. $|\alpha'_3| \leq |k|^4$ for $|k| \geq 10^7$.

□

7.3. The final result.

We denote Mahler measure as $M(\alpha)$ and logarithmic Weil's height as $h(\alpha)$.

Lemma 7.3. *If $|k| \geq 5 \cdot 10^{37}$, $\Gamma \neq 0$ and $V_n = W_m$, then $m \leq 2$ or $n \leq 2$.*

Proof. Assume that, on the contrary, $V_n = W_m$ and $n \geq m \geq 3$. It holds that $M(\alpha_3) \leq |a_d| \prod_{i=1}^d \max\{|\alpha'_i|, 1\}$ and $|a_d| \leq \left(\sqrt{|k|+1}(|x_1| \sqrt{16|k|^3+4|k|} + |z_1| \sqrt{|k|+1}) \right)^{32} < 257^{16}|k|^{65}$. Since Lemma 7.2 provides the bound for conjugates $|\alpha'_3| \leq |k|^4$, it follows that

$$h(\alpha_3) \leq \frac{1}{32} \log(257^{16}|k|^{65} \cdot |k|^{4 \cdot 32}) = 2.774538 + \frac{193}{32} \log |k|.$$

For all three bounds it holds that $h'(\alpha_i) \leq 7 \log |k|$.

Lemma 2.6 implies $|\Gamma| = |\log \Lambda'| < K \sqrt{|ac|} |s + \sqrt{ac}|^{-m}$ (if $m, n \geq 3$), where $K = \frac{8}{3} \log \frac{24}{19} = 0.622973$. Since $|s + \sqrt{ac}| \geq \sqrt{|ac|} = \sqrt{|(16k^3 - 4k)(k - 1)|}$, it follows that $|s + \sqrt{ac}| > 3|k|^2$ (for $|k| > 3$). Hence

$$|\Gamma| < K \sqrt{|ac|} (3|k|^2)^{-m} < K(1.5|k|)^{1-m}.$$

We now apply the following well-known theorem from [4].

Theorem 7.4 (Baker, Wüstholz). *Let $\Gamma = b_1 \log \alpha_1 + b_2 \log \alpha_2 + \dots + b_n \log \alpha_n$ be a linear form in logarithms of algebraic numbers $\alpha_1, \alpha_2, \dots, \alpha_n$ with integer coefficients b_1, b_2, \dots, b_n . If $\Gamma \neq 0$, then*

$$\log |\Gamma| \geq -18(n+1)! n^{n+1} (32d)^{n+2} \log(2nd) h'(\alpha_1) h'(\alpha_2) \dots h'(\alpha_n) \log B,$$

where $d = [\mathbb{Q}(\alpha_1, \alpha_2, \dots, \alpha_n) : \mathbb{Q}]$, $B = \max\{|b_1|, |b_2|, \dots, |b_n|\}$, and $h'(\alpha) = \max\{h(\alpha), \frac{1}{d} |\log \alpha|, \frac{1}{d}\}$.

We use the logarithm only for absolute values (positive reals), but this theorem holds more generally. For the logarithm of a complex number $z = re^{i\varphi}$ with $r > 0$ we can take $\log z = \log r + i\varphi$.

Baker–Wüstholz theorem implies that, if $\Gamma \neq 0$, then

$$\log K(1.5|k|)^{1-m} > \log |\Gamma| > -18 \cdot 4! \cdot 3^4 (32 \cdot 2048)^5 \cdot 343 \log^3 |k| \log(6 \cdot 2048) \log n,$$

meaning $(1-m) \log \frac{3}{2}|k| > \log K + (1-m) \log \frac{3}{2}|k| > -K' \log^3 |k| \log n$, where $\log K \approx -0.47325$. Together with Lemma 7.1, this implies that

$$\frac{m-1}{\log(3m+2)} \leq \frac{m-1}{\log n} < K' \frac{\log^3 |k|}{\log \frac{3}{2}|k|} < K' \log^2 |k|,$$

where $K' = 18 \cdot 4! \cdot 3^4 (32 \cdot 2048)^5 \cdot 343 \log(6 \cdot 2048) \approx 1.3663 \cdot 10^{32}$.

Since $m \geq 2|k| - 1$ (by (5.1)) and since function $f(x) = \frac{x-1}{\log(3x+2)}$ is increasing, it follows that $\frac{2|k| - 1}{\log(6|k| - 1)} < K' \log^2 |k|$ and $|k| - 1 < 6.831506 \cdot 10^{31} \log^2 |k| \log(6|k| - 1)$, which is impossible for $|k| \geq 5 \cdot 10^{37}$. \square

Lemma 7.5. *If k is a Gaussian integer such that $\operatorname{Im} k \operatorname{Re} k \neq 0$, then $\frac{|k+1|}{|k-1|} \notin \mathbb{Q}$.*

Proof. It is sufficient to show that $|k+1| \cdot |k-1|$ is not in \mathbb{Q} , so it also suffices to show that it is not a rational integer. If $k = x + yi$, then $|k+1|^2 \cdot |k-1|^2 = x^4 + y^4 + 1 + 2x^2y^2 - 2x^2 + 2y^2$, so we need to show that this expression is not a perfect square.

It holds that $x^4 + y^4 + 1 + 2x^2y^2 - 2x^2 + 2y^2 > (x^2 + y^2 - 2)^2$, because this is equivalent to $2x^2 + 6y^2 > 3$. Similarly, $x \neq 0$ implies that $x^4 + y^4 + 1 + 2x^2y^2 - 2x^2 + 2y^2 < (x^2 + y^2 + 1)^2$.

From $x^4 + y^4 + 1 + 2x^2y^2 - 2x^2 + 2y^2 = (x^2 + y^2)^2$, it follows that $1 - 2x^2 + 2y^2 = 0$, which is impossible (parity check), while $x^4 + y^4 + 1 + 2x^2y^2 - 2x^2 + 2y^2 = (x^2 + y^2 - 1)^2$ implies that $4y^2 = 0$, again impossible since $y \neq 0$. \square

Lemma 7.6. *If k is a Gaussian integer such that $\operatorname{Im} k \neq 0$, then $\frac{|16k^3 - 4k|}{|k-1|} \notin \mathbb{Q}$.*

Proof. Again, it suffices to show that $|16k^3 - 4k| \cdot |k-1| \notin \mathbb{Z}$, i. e. $|4k^3 - k|^2 \cdot |k-1|^2$ is not a perfect square.

If $k = x + yi$, then $|4k^3 - k|^2 \cdot |k-1|^2 = |4x^4 - 4x^3 - 24x^2y^2 - x^2 + 12xy^2 + x + 4y^4 + y^2 + i(16x^3y - 12x^2y - 16xy^3 - 2xy + 4y^3 + y)|^2 = (x^2 - 2x + 1 + y^2)(4x^2 - 4x + 1 + 4y^2)(4x^2 + 4x + 1 + 4y^2)(x^2 + y^2)$. By substituting $z = -4x + 1$, we get $(z^4 + (32y^2 - 10)z^2 + 160y^2 + 256y^4 + 9)^2 + 4096y^2z^2$. Further substitutions $u = z^2, v = y^2$ give that $(u^2 + (32v - 10)u + 160v + 256v^2 + 9)^2 + 4096uv$ is a square, where u and v are also perfect squares. Since $y = \operatorname{Im} k \neq 0$, we see that $v \neq 0$, and neither $u = (-4x + 1)^2 = 0$. Hence $(u^2 + (32v - 10)u + 160v + 256v^2 + 9)^2 + 4096uv > (u^2 + (32v - 10)u + 160v + 256v^2 + 9)^2$. If the left-hand side is a square, there is a positive integer w such that $(u^2 + (32v - 10)u + 160v + 256v^2 + 9)^2 + 4096uv = (u^2 + (32v - 10)u + 160v + 256v^2 + 9 + w)^2$.

The equation $2w(u^2 + (32v - 10)u + 160v + 256v^2 + 9) + w^2 - 4096uv = 0$ is quadratic in u . The discriminant is

$$\begin{aligned} 4D(v, w) &= -8(w^3 - 32w^2 + 65536v^2w - 2097152v^2 + 640vw^2 - 20480vw) \\ &= -4 \cdot 2((w^2(w - 32) + 65536v^2(w - 32) + 640wv(w - 32)), \end{aligned}$$

which is negative for $w > 32$. For a solution to be an integer, the discriminant must be a perfect square. It follows that $w \in \{1, 2, \dots, 32\}$.

Since $D(v, 32) = 0$, solving the quadratic equation implies that $u = 16v + 5$, which is not a square because 5 is not a quadratic remainder modulo 16. For the most of the remaining values of w , in a similar manner we show that $D(v, w)$ is not a square. Observe that $D(v, w) \equiv -2w^2(w - 32) \pmod{128}$.

For odd w , we get that $D(v, w) \equiv 2 \pmod{4}$, which cannot be a square. For $w \equiv 2 \pmod{8}$, it holds that $D(v, w) \equiv -16 \pmod{64}$, while for $w \equiv 4 \pmod{8}$, $D(v, w) \equiv 128 \pmod{256}$ and again the discriminant cannot be a square.

For $w = 6$, $\frac{D(v,6)}{16} = 212992v^2 + 12480v + 117 \equiv 5 \pmod{16}$, so $D(v, 6)$ is not a square. Similarly, $\frac{D(v,8)}{256} \equiv 12 \pmod{16}$, $\frac{D(v,16)}{4096} \equiv 2 \pmod{4}$ and $\frac{D(v,22)}{16} \equiv 13 \pmod{16}$ imply that none of $D(v, 8)$, $D(v, 16)$ and $D(v, 22)$ can be a square.

For $w = 14$, it holds that $(128v + 9)^2 > \frac{D(v,14)}{144} = (128v + 7)^2 + 448v$, hence $D(v, 14)$ is not a square since $v > 0$. Similarly, $(32v + 4)^2 > \frac{D(v,24)}{1024} = 1024v^2 + 240v + 9 = (32v + 3)^2 + 48$ and $(128v + 19)^2 > \frac{D(v,30)}{16} = (128v + 15)^2 + 960v$ show that neither $D(v, 24)$ nor $D(v, 30)$ is a square ($v > 0$).

We have checked all $w \in \{1, 2, \dots, 31, 32\}$ and thus proven the lemma. \square

Theorem 7.7. *Let k be a Gaussian integer such that $\operatorname{Re} k \neq 0$ and $|k| \geq 5 \cdot 10^{37}$. The Diophantine triple $\{k - 1, k + 1, 16k^3 - 4k\}$ can be extended to a Diophantine quadruple only by $d = 4k$ or $d = 64k^5 - 48k^3 + 8k$.*

Proof. If $\operatorname{Im} k = 0$, then the elements of the sequence $(V_n)_n$ are integers, hence x is an integer too. Since $(k - 1)d + 1 = x^2$, it follows that $d \in \mathbb{Q} \cap \mathbb{Z}[i]$, so d is an integer. The sign of d is the same as the sign of $k - 1, k + 1$ and $16k^3 - 4k$ (because $d = \frac{x^2 - 1}{k - 1} \neq 0$), so Theorem 1 from [6], since $|k| \geq 2$, implies that $d = 4k$ or $d = 64k^5 - 48k^3 + 8k$.

From now on, we assume that $\{a, b, c, d\}$ is a Diophantine quadruple for $a = k - 1, b = k + 1, c = 16k^3 - 4k$ and that $\operatorname{Im} k$ is not 0.

Checking V_n and W_m for small indices, we obtain the extensions $4k, 64k^5 - 48k^3 + 8k$ and candidates such as $W_1^{(1)} = 4k^2 - k - 2$. By computing the first few elements of $(V_n)_n$, $V_1 = 2k - 1$ and $V_2 = 4k^2 - 2k - 1$, we see that $W_1^{(2)}$ cannot have the same value (for large $|k|$), and neither can the larger elements V_n for $n \geq 3$, since these are greater in absolute value than $W_1^{(1)}$: $|V_n| - |W_1^{(1)}| \geq |V_3 - W_1^{(1)}| = |8k^3 - 4k^2 - 4k + 1 - (4k^2 - k - 2)| > 0$ for $|k| > 10^{37}$. Similarly, $V_1 = 2k - 1$ and $V_2 = 4k^2 - 2k - 1$ cannot be an element of the sequence $W_m^{(1)}$. Analogously we check sequences $W_m^{(j)}$ for the remaining $j = 2, 3, 4, 5, 6$.

Therefore, indices n and m are greater than 2 if $d \notin \{4k, 64k^5 - 48k^3 + 8k\}$.

Lemma 7.5 and Lemma 7.6 imply that $\frac{|c|}{|a|}$ and $\frac{|b|}{|a|}$ are not rational numbers. In the same manner as in the Lemma 3.1, this implies that the linear form $\Gamma = \log \Lambda'$ is not 0. If $V_n = W_m$ for $m \geq 2$ and $n \geq 2$, then Lemma 7.3 would imply that $|k| < 5 \cdot 10^{37}$, which is a contradiction. Therefore, the assumption that $V_n = W_m$ for $m \geq 3$ and $n \geq 3$ is wrong, and so is the claim that $d \notin \{4k, 64k^5 - 48k^3 + 8k\}$ for $|k| \geq 5 \cdot 10^{37}$. \square

ACKNOWLEDGEMENTS

N. A. and A. F. were supported by the Croatian Science Foundation under the project no. IP-2018-01-1313.

REFERENCES

1. N. Adžaga, *On the size of Diophantine m -tuples in imaginary quadratic number rings*, Bulletin of Mathematical Sciences (to appear).
2. Lj. Bačić, *Sets in which $xy + 4$ is always a square and problem of the extensibility of some parametric Diophantine triples*, PhD Thesis (2014), Faculty of Science, University of Zagreb.
3. A. Baker and H. Davenport, *The equations $3x^2 - 2 = y^2$ and $8x^2 - 7 = z^2$* , Quarterly Journal of Mathematics. Oxford. Second Series **20** (1969), no. 2, 129–137.
4. A. Baker and G. Wüstholz, *Logarithmic forms and group varieties*, Journal für die reine und angewandte Mathematik **442** (1993), 19–62.
5. A. Bayad, A. Filipin, and A. Togbé, *Extension of a parametric family of Diophantine triples in Gaussian integers*, Acta Mathematica Hungarica **148** (2016), no. 2, 312–327.
6. Y. Bugeaud, A. Dujella, and M. Mignotte, *On the family of Diophantine triples $\{k - 1, k + 1, 16k^3 - 4k\}$* , Glasgow Mathematical Journal **49** (2007), 333–344.
7. A. Dujella, *An absolute bound for the size of Diophantine m -tuples*, Journal of Number Theory **89** (2001), 126–150.
8. L. Fjellstedt, *On a class of Diophantine equations of second degree in imaginary quadratic fields*, Arkiv för Matematik **2** (1953), no. 24, 435–461.
9. Z. Franušić, *On the extensibility of Diophantine triples $\{k - 1, k + 1, 4k\}$ for Gaussian integers*, Glasnik matematički **43** (2008), no. 2, 265–291.
10. Z. Franušić and B. Jadrijević, *Computing relative power integral bases in a family of quartic extensions of imaginary quadratic fields*, Publicationes Mathematicae Debrecen **92** (2018), 293–315.
11. B. He, A. Togbé, and V. Ziegler, *There is no Diophantine quintuple*, Transactions of the American Mathematical Society **371** (2019), no. 9, 6665–6709.
12. B. Jadrijević and V. Ziegler, *A system of relative Pellian equations and a related family of relative Thue equations*, International Journal of Number Theory **2** (2006), no. 4, 569–590.
13. N. P. Smart, *The algorithmic resolution of Diophantine equations*, London Mathematical Society, Cambridge University Press, 1998.
14. Wolfram Research Inc., *Mathematica*, 2018, Version 11.3.0, Champaign, Illinois, USA.

NIKOLA ADŽAGA, FACULTY OF CIVIL ENGINEERING, UNIVERSITY OF ZAGREB
 E-mail address: nadzaga@grad.hr

ALAN FILIPIN, FACULTY OF CIVIL ENGINEERING, UNIVERSITY OF ZAGREB
 E-mail address: filipin@grad.hr

ZRINKA FRANUŠIĆ, FACULTY OF SCIENCE, UNIVERSITY OF ZAGREB
 E-mail address: fran@math.hr