

# q-spherical surfaces in Euclidean space

---

**Gorjanc, Sonja; Jurkin, Ema**

*Source / Izvornik:* **Filomat, 2023, 37, 1 - 11**

**Journal article, Published version**

**Rad u časopisu, Objavljena verzija rada (izdavačev PDF)**

<https://doi.org/10.2298/FIL2301001G>

*Permanent link / Trajna poveznica:* <https://um.nsk.hr/um:nbn:hr:237:358960>

*Rights / Prava:* [In copyright](#)/[Zaštićeno autorskim pravom.](#)

*Download date / Datum preuzimanja:* **2025-03-29**

*Repository / Repozitorij:*

[Repository of the Faculty of Civil Engineering,  
University of Zagreb](#)





## $q$ -spherical surfaces in Euclidean space

Sonja Gorjanc<sup>a</sup>, Ema Jurkin<sup>b</sup>

<sup>a</sup>University of Zagreb, Faculty of Civil Engineering

<sup>b</sup>University of Zagreb, Faculty of Mining, Geology and Petroleum Engineering

**Abstract.** In this paper we define  $q$ -spherical surfaces as the surfaces that contain the absolute conic of the Euclidean space as a  $q$ -fold curve. Particular attention is paid to the surfaces with singular points of the highest order. Two classes of such surfaces, with one and two  $n$ -fold points, are discussed in detail. We study their properties, give their algebraic equations and visualize them with the program *Mathematica*.

### 1. Motivation

One of the most important places in the classical geometry belongs to the study of some classes of surfaces with special properties in the Euclidean space. In the real projective plane the Euclidean metric defines the Euclidean plane  $\mathbb{E}^2$  with the *absolute (circular) points*  $(0, 1, i)$  and  $(0, 1, -i)$  (i.e.  $x_0 = 0, x_1^2 + x_2^2 = 0$ ). The absolute points do not belong to the real plane, they belong to its complexification. In this paper we consider both real and imaginary elements. An algebraic curve of order  $n$  passing through the absolute points is called *circular curve*. If it contains absolute points as its  $q$ -fold points, the curve is called  *$q$ -circular*, and if  $n = 2q$  the curve is called *entirely circular*. Some well-known circular curves are strophoid, trisectrix of Maclaurin, limaçon, cardioid, lemniscate of Bernoulli, Booth lemniscate, Cartesian ovals, Cassini ovals, astroid and Watt's curve. Due to their numerous applications in engineering, the circular curves in the Euclidean plane have been treated in a significant number of papers (see e.g. [1], [3], [19], [20], [21], [22]). On the other hand, the surfaces in the Euclidean space with the similar properties, with an exception of the cyclides that will be mentioned later, have been in some way neglected. Motivated by this fact, we introduce so-called  $q$ -spherical surfaces, the surfaces that contain the absolute conic of  $\mathbb{E}^3$  as a  $q$ -fold curve. We study some of their properties and visualize their shapes. Particular attention is paid to the surfaces with singular points of the highest order.

### 2. Introduction

In the real three-dimensional projective space  $P^3(\mathbb{R})$ , in homogeneous Cartesian coordinates  $(x_0, x_1, x_2, x_3)$ , ( $x_1, x_2, x_3 \in \mathbb{R}, x_0 \in \{0, 1\}, (x_0, x_1, x_2, x_3) \neq (0, 0, 0, 0)$ ), the equation

$$F_n(x_0, x_1, x_2, x_3) = 0,$$

---

2020 *Mathematics Subject Classification*. Primary 51N20; Secondary 51M15.

*Keywords*.  $q$ -spherical surface; Absolute conic; Singular point.

Received: 19 May 2020; Accepted: 02 February 2022

Communicated by Dragan S. Djordjević

*Email addresses*: sgorjanc@grad.hr (Sonja Gorjanc), ema.jurkin@rgn.hr (Ema Jurkin)

where  $F_n$  is a homogeneous polynomial of degree  $n$ , defines an  $n^{\text{th}}$  order surface  $S_n$ . This equation can be written as

$$f_n(x_1, x_2, x_3) + x_0 f_{n-1}(x_1, x_2, x_3) + \dots + x_0^{n-1} f_1(x_1, x_2, x_3) + x_0^n f_0(x_1, x_2, x_3) = 0, \quad (1)$$

where  $f_j$ ,  $j = 1, \dots, n$ , are homogeneous polynomials of degree  $j$ , [16].

Any straight line, not lying on  $S_n$ , intersects  $S_n$  in  $n$  points and any plane intersects  $S_n$  in the  $n^{\text{th}}$  order plane curve.

A point  $T$  of the surface  $S_n$  for which at least one partial derivative of  $F_n$  is not equal to zero is called a *regular point* of  $S_n$ . All tangents to the surface at that point lie in one plane - the tangent plane of  $S_n$  at  $T$ .

A point  $T$  of the surface  $S_n$  for which all partial derivatives of  $F_n$  are equal to zero is called a *singular point* of  $S_n$ . The tangents to  $S_n$  at this point form an algebraic cone with vertex in  $T$ . If the tangent cone is of order  $k$ , the point  $T$  is the *k-fold point* of the surface  $S_n$ . Every plane through  $T$  intersects  $S_n$  in the  $n^{\text{th}}$  order plane curve with the  $k$ -fold point in  $T$ .

The point  $T$  is a  $k$ -fold point of  $S_n$  if all partial derivatives of  $F_n$  with the order less than  $k$  vanish at  $T$ , and at least one  $k$ -order derivative of  $F_n$  at  $T$  doesn't vanish, [4].

According to [10], if the origin  $O(1, 0, 0, 0)$  is the  $k$ -fold point of  $S_n$ , then  $S_n$  has the equation,

$$f_n(x_1, x_2, x_3) + x_0 f_{n-1}(x_1, x_2, x_3) + \dots + x_0^{n-k} f_k(x_1, x_2, x_3) = 0, \quad (2)$$

and the tangent cone at  $O$  is given by

$$f_k(x_1, x_2, x_3) = 0. \quad (3)$$

If every point of a curve  $C$  lying on the surface  $S_n$  is the  $k$ -fold point of  $S_n$ , then  $C$  is a *k-fold curve* of  $S_n$ .

In the real projective space  $P^3(\mathbb{R})$  the Euclidean metric defines the Euclidean space  $\mathbb{E}^3$  with the absolute conic given by the equations:

$$x_0 = 0 \text{ and } A_2 = x_1^2 + x_2^2 + x_3^2 = 0.$$

The absolute conic does not belong to the real Euclidean space, but to its complexification.

### 3. $q$ -spherical surfaces

**Definition 3.1.** A surface  $S_n$  of Euclidean space is called  $q$ -spherical surface if it contains the absolute conic as a  $q$ -fold curve.

**Theorem 3.2.** In Euclidean space  $q$ -spherical surface of order  $n$  is given by the following equation:

$$A_2^q g_{n-2q}(x_1, x_2, x_3) + \sum_{j=1}^{q-1} x_0^j A_2^{q-j} g_{n-2q+j}(x_1, x_2, x_3) + \sum_{j=q}^n x_0^j f_{n-j}(x_1, x_2, x_3) = 0, \quad (4)$$

where  $n \geq 2q$ ,  $g_{n-2q} \neq 0$ ,  $A_2 \nmid g_{n-2q}$ ,  $f_{n-q} \neq 0$  and  $A_2 \nmid f_{n-q}$ .

**PROOF:** Let us first prove that the surface given by (4) is  $q$ -spherical surface. For all points on the absolute conic ( $A_2 = 0$ ,  $x_0 = 0$ ) all first  $q - 1$  partial derivatives of the polynomial on the left hand side of (4) vanish since their terms contain either  $A_2$  or  $x_0$  as a factor. According the theorem's condition, at least one  $q$ -order derivative does not vanish for the points of the absolute conic.

Now, let a  $q$ -spherical surface of order  $n$  be given by the equation (1) that can be written in the following form:

$$F_n(x_0, x_1, x_2, x_3) = f_n(x_1, x_2, x_3) + \sum_{j=1}^{q-1} x_0^j f_{n-j}(x_1, x_2, x_3) + \sum_{j=q}^n x_0^j f_{n-j}(x_1, x_2, x_3) = 0. \quad (5)$$

Since the first  $q - 1$  partial derivatives of the polynomial  $F_n$  have to vanish for  $A_2 = 0$  and  $x_0 = 0$ , it is clear that the polynomials  $f_{n-j}$ ,  $j = 1, \dots, q - 1$ , must contain  $A_2^{q-j}$  as a factor. Since at least one  $q$ -order derivative of  $F_n$  doesn't vanish, it is clear that  $f_{n-q} \neq 0$  and  $A_2$  is not a factor of  $f_{n-q}$ .  $\square$

In [9] and [5] the authors studied the special class of 1-spherical surfaces, the surfaces which touch the plane at infinity through the absolute conic and have a singular point of the highest order. These surfaces belong to a wider class of surfaces, so-called monoid surfaces, also treated in [12] and introduced in [15].

In paper [7] the authors considered a congruence of circles  $C(p)$  that consists of circles in Euclidean space  $\mathbb{E}^3$  passing through two given points  $P_{1,2}(0, 0, \pm p)$ . For a given congruence  $C(p)$  and a given curve  $\alpha$ , a circular surface  $CS(\alpha, p)$  is defined as the system of circles from  $C(p)$  that intersect  $\alpha$ . If  $\alpha$  is an  $m^{\text{th}}$  order algebraic curve that intersects the axis  $z$  at  $z'$  points, the absolute conic at  $a'$  pairs of the absolute points and with the points  $P_1$  and  $P_2$  as  $p'_1$ -fold and  $p'_2$ -fold points, respectively, then, the following statements hold:

–  $CS(\alpha, p)$  is an algebraic surface of the order  $3m - (z' + 2a' + 2p'_1 + 2p'_2)$ .

– The absolute conic is an  $m - (z' + p'_1 + p'_2)$ -fold curve of  $CS(\alpha, p)$ .

– The axis  $z$  is an  $(m - 2a' + z')$ -fold line of  $CS(\alpha, p)$ .

– The points  $P_1, P_2$  are  $2m - (2a' + p'_1 + p'_2)$ -fold points of  $CS(\alpha, p)$ . (CS)

Thus, the most of these surfaces are spherical, and some of them will be studied in more detail in subsection 4.1.2.

The further examples of  $q$ -spherical surfaces can be found in [6] and [8] where authors introduced rose surfaces and generalized rose surfaces as the special cases of circular surfaces for which  $\alpha$  is a cyclic-harmonic curve. It was shown how  $q$  depends on the properties and position of the referent cyclic-harmonic curve with respect to the singular points of the congruence  $C(p)$ .

#### 4. Entirely spherical surfaces

**Definition 4.1.** A surface  $S_{2n}$  of Euclidean space is called entirely spherical surface if it contains the absolute conic as an  $n$ -fold curve.

Probably the mostly studied surfaces of the fourth order are cyclides, the bispherical quartic surfaces, [17].

The term "cyclides" is often used for their special class, Dupin cyclides, which can be defined in a several ways, [2], [11]. They are only surfaces that have the property that their evolulute degenerate into curves, in fact both sheets of the focal surface are conics. The examples of cyclides are the tori, cones and cylinders of revolutions. Dupin cyclides can also be defined as the envelopes of the family of spheres tangent to three fixed spheres. Dupin cyclides are the inverse images of the standard tori, cylinders or cones. The inverse image of a ring torus, horn torus and spindle torus are called a ring cyclide, horn cyclide and spindle cyclide, respectively. If the center of the inversion sphere lies on the torus, the obtained surface is a parabolic cyclide (ring, horn or spindle), [23]. The cyclides as the surfaces of the fourth order having the circle at infinity as the nodal conic were studied in [17]. Their equation of type (4) for  $n = 4, p = 2$ , was given. The cyclide was also defined as the envelope of sphere whose center moves on a fixed quadric, and which intersects a fixed sphere orthogonally. The intersection curve of the fixed quadric and sphere is the focal curve of the cyclide. The number of strait lines lying on the cyclide is sixteen. In [17] the author distinguishes 23 types of cyclides.

In [14] the family of surfaces called Darboux cyclides were studied. These surfaces are algebraic surfaces of order at most 4 and are a superset of bispherical surfaces of order 4, circular surfaces of order 3 and quadrics, and they carry up to six real families of circles.

Entirely spherical surfaces can be constructed in many different ways. Let us mention some of them:

An inverse image of a surface  $S_n$  of order  $n$  is an  $n$ -spherical surface  $S_{2n}$  of order  $2n$  with an  $n$ -fold point in the pole of inversion.

If  $C_n$  is a surface of class  $n$  and  $P$  a point in general position to  $C_n$ , then the pedal surface of  $C_n$  with respect to the pole  $P$  is an  $n$ -spherical surface  $S_{2n}$  of order  $2n$  with an  $n$ -fold point in the pole  $P$ , [13].

According to the properties (CS) of circular surfaces  $\mathcal{CS}(\alpha, p)$  that are pointed out in the previous section, if  $\alpha = k_{2n}$  is an entirely circular curve of order  $2n$  with an  $n$ -fold point in  $P_1$  ( $m = 2n$ ,  $a' = p_1 = n$ ,  $p_2 = 0$ ,  $z' = 0$ ), the obtained surface  $\mathcal{CS}(k_{2n}, p)$  is entirely spherical surface with two  $n$ -fold points  $P_1, P_2$ . The examples of these surfaces will be given in subsection 4.1.2.

**Theorem 4.2.** *An entirely spherical surface  $S_{2n}$  of order  $2n$  can't have singular points of multiplicity higher than  $n$ .*

PROOF: If there was a point  $T$  of multiplicity  $n + 1$ , all isotropic lines through  $T$  would lie on  $S_{2n}$  and the surface would split onto the isotropic cone with vertex in  $T$  and a surface of order  $2n - 2$ .  $\square$

**Lemma 4.3.** *A plane curve of order  $2n$ ,  $n \geq 2$ , can have at most three  $n$ -fold points.*

PROOF: The maximum number of double points of a curve of order  $2n$  equals  $D = \frac{(2n-1)(2n-2)}{2}$ , [18]. Every  $n$ -fold point is counted as  $N = \frac{n(n-1)}{2}$  double points. Since  $\frac{D}{N} = 4 - \frac{2}{n}$ , the curve of order  $2n$  can have three, but not four  $n$ -fold points.  $\square$

Therefore, if a plane curve of order  $2n$  has more than three  $n$ -fold points, it splits onto the curves of lower order.

**Lemma 4.4.** *An entirely circular curve of order  $2n$ ,  $n \geq 2$ , can have only one  $n$ -fold point beside the absolute points.*

PROOF: The statement follows directly from Lemma 4.3.  $\square$

**Theorem 4.5.** *If an entirely spherical surface  $S_{2n}$  contains two real and distinct  $n$ -fold points  $N_1$  and  $N_2$ , every plane through the line  $N_1N_2$  intersects  $S_{2n}$  along  $n$  circles passing through  $N_1$  and  $N_2$ .*

PROOF: Let  $N_1$  and  $N_2$  be the  $n$ -fold points of the surface  $S_{2n}$ , and let  $\beta$  be a plane through  $N_1$  and  $N_2$ . The intersection line of  $\beta$  and  $S_{2n}$  is a curve  $k_{2n}$  of order  $2n$ . According to Lemma 4.4 this curve is not proper, it splits onto curves  $k_{2t}$  and  $k_{2n-2t}$  of order  $2t$  and  $2n - 2t$ , respectively. Since  $k_{2n}$  passes through the absolute points  $n$  times, the curves  $k_{2t}$ ,  $k_{2n-2t}$  have to be entirely circular passing through the absolute points  $t$  and  $n - t$  times. The necessary multiplicity of the point  $N_1$  will be achieved if it is the singular point of  $k_{2t}$  and  $k_{2n-2t}$  with multiplicity  $t$  and  $n - t$ . Now, again according to Lemma 4.4,  $N_2$  is the singular point of  $k_{2t}$  and  $k_{2n-2t}$  with multiplicity at most  $t - 1$  and  $n - t - 1$ . This is in contradiction with the fact that  $k_{2n}$  passes  $n$  times through  $N_2$ . It follows that  $k_{2t}$  and  $k_{2n-2t}$  are not proper curves either. Therefore, they again split onto the curves of lower order. Continuing with this procedure, in the last step we come to the  $n$  curves of order 2, i.e. circles.  $\square$

#### 4.1. Examples

In this subsection we give some examples of the spherical surfaces. For a particular surface given by an implicit equation, we study its properties and visualize its shape. For computing and plotting, we use the program *Mathematica*. In some examples we start with an algebraic equation of the surface and determine its properties. In other cases we first give a construction of the surface from which we derive its equation.

##### 4.1.1. Entirely spherical surfaces $S_{2n}$ with only one $n$ -fold point

A surface  $S_{2n}$  given by the equation of the form

$$A_2(x_1, x_2, x_3)^n + x_0^n \cdot f_n(x_1, x_2, x_3) = 0,$$

i.e. in the affine coordinates

$$A_2(x, y, z)^n + f_n(x, y, z) = 0, \tag{6}$$

is an entirely spherical surface with the  $n$ -fold point at the origin.

**Proposition 4.6.** *There are only  $2n$  straight lines through the origin lying entirely on the surface  $S_{2n}$  given by (6). These lines are the intersections of cones given by*

$$A_2(x, y, z) = 0, \quad f_n(x, y, z) = 0,$$

and they are imaginary in pairs.

**PROOF:** Let a line  $p$  through  $O(0,0,0)$  be spanned by  $O$  and a further point  $P(a, b, c) \neq O$ . The line  $p$  is parametrized by

$$p \quad \dots \quad (x, y, z) = (ta, tb, tc), t \in \mathbb{R}. \tag{7}$$

It lies on  $S_{2n}$  if and only if

$$A_2(ta, tb, tc)^n + f_n(ta, tb, tc) = 0,$$

for every  $t \in \mathbb{R}$ . This is precisely when

$$t^n [t^n A_2(a, b, c)^n + f_n(a, b, c)] = 0,$$

for every  $t \in \mathbb{R}$ . It follows that  $A_2(a, b, c) = 0$  and  $f_n(a, b, c) = 0$ . Therefore,  $A_2(ta, tb, tc) = 0$ ,  $f_n(ta, tb, tc) = 0$ , for every  $t \in \mathbb{R}$ . Evidently the line  $p$  lies on the cones given by equations  $A_2(x, y, z) = 0$  and  $f_n(x, y, z) = 0$ . We conclude: the only lines through the origin that lie on  $S_{2n}$  are the isotropic lines on the tangent cone at the origin.  $\square$

**Proposition 4.7.** *The surface  $S_{2n}$  given by (6) has no other singular points beside the origin.*

**PROOF:** This fact can be proved as follows: Let a non-isotropic line  $p$  through the origin be given by (7). We compute the intersections of  $S_{2n}$  and  $p$ . They belong to the zeros of the following polynomial of degree  $2n$  in  $t$ :

$$P(t) := A_2(ta, tb, tc)^n + f_n(ta, tb, tc).$$

Obviously,  $t = 0$  is zero with multiplicity  $n$ . The other  $n$  zeros are given by

$$t^n = -\frac{f_n(a, b, c)}{(a^2 + b^2 + c^2)^n}$$

and therefore different (in general) complex numbers.  $\square$

The tangent cone  $\Phi_n$  at the  $n$ -fold point  $O$  of the surface  $S_{2n}$  given by (6) has the equation  $f_n(x, y, z) = 0$ . The polynomial  $f_n$  can be irreducible or reducible. If the polynomial  $f_n$  can be factorized

$$f_n(x, y, z) = f_{n_1}(x, y, z) \cdot \dots \cdot f_{n_k}(x, y, z), \quad n_1 + \dots + n_k = n, \tag{8}$$

the tangent cone  $\Phi_n$  splits into the cones  $\Phi_{n_1}, \dots, \Phi_{n_k}$  of order  $n_1, \dots, n_k$ , respectively. Therefore, the classification of surfaces  $S_{2n}$  can be made according to the degrees of the polynomials  $f_{n_1}, \dots, f_{n_k}$ . If we assume that all polynomials  $f_{n_1}, \dots, f_{n_k}$  determine different cones  $\Phi_{n_1}, \dots, \Phi_{n_k}$ , then for all  $n \in \mathbb{N}$  the surfaces  $S_{2n}$  can be classified into  $p(n)$  types, where  $p$  is the partition function, i.e.  $p(n)$  is the number of ways of writing the integer  $n$  as a sum of positive integers, where the order of addends is not considered significant. The formulas for counting the value  $p(n)$  can be found in [24].

Some examples of surfaces given by (6), together with their tangent cones at the  $n$ -fold points, are shown in Figures 1-4.




Figure 1: Five examples for  $n = 2, 3, \dots, 6$  where the tangent cone at the origin splits into  $n$  planes, i.e.  $f_n(x, y, z) = \prod_{i=1}^n h_i(x, y, z)$ , where  $\forall i, h_i(x, y, z)$  is a linear polynomial.




Figure 2: Surfaces  $S_{2n}$  with the proper tangent cones of degree  $n$  at the origin, where  $f_n(x, y, z)$  is:  $x^2 + y^2 - z^2$  (a),  $-(x^2 + y^2 - z^2)$  (b),  $-2x^3 - x^2z + 2y^2z + xz^2$  (c),  $48x^4 + 48y^4 - 64\sqrt{3}y^3z + 40y^2z^2 - z^4 + 8x^2(12y^2 + 24\sqrt{3}yz + 5z^2)$  (d).




Figure 3: Three examples for  $n = 3$  where the tangent cone at the origin splits into a plane and  $2^{\text{nd}}$  degree cone.




Figure 4: Four examples for  $n = 4$  where the tangent cone at the origin splits into two  $2^{\text{nd}}$  degree cone or into two planes and one  $2^{\text{nd}}$  degree cone.

4.1.2. Entirely spherical surfaces  $S_{2n}$  with two  $n$ -fold points

In this subsection we study a class of entirely spherical surfaces having two singular points of the highest order. We start with defining a class of entirely circular curves.

A curve  $k^{2n}$  of order  $2n$ ,  $n \geq 2$ , given by the equation

$$(x^2 + y^2)^n + f_n(x, y) = 0, \tag{9}$$

where homogeneous polynomial  $f_n$  is a product of  $n$  linear factors, is an entirely circular curve having an  $n$ -fold point at the origin. Linear factors of  $f_n$  represent the tangent lines at the singular point. If the tangent lines divide the plane into equal parts, polynomial  $f_n$  equals

$$f_n = \begin{cases} \prod_{i=0}^{n-1} \left( \cos i \frac{2\pi}{n} \cdot y - \sin i \frac{2\pi}{n} \cdot x \right), & n \text{ odd,} \\ \prod_{i=0}^{n-1} \left( \cos i \frac{\pi}{n} \cdot y - \sin i \frac{\pi}{n} \cdot x \right), & n \text{ even.} \end{cases}$$



Some examples of this type of curves are shown in Figure 5.




Figure 5: Five examples of entirely circular curves with equation (9) for  $n = 2, \dots, 6$ .

**Remark 4.8.** The examples in Figure 5 make it easy to see that every straight line through the origin intersects  $k^{2n}$ , except at the  $n$ -fold point  $O$ , at just one real point if  $n$  is odd number, and that the number of real intersections is zero or two if  $n$  is even. This fact can be also numerically verify for any chosen  $n$  and straight line through  $O$  using the program Mathematica.

Switching to polar coordinates  $O(\rho, \varphi)$ , where  $x = \rho \cos \varphi$  and  $y = \rho \sin \varphi$ , we obtain the following polar equation of the curve  $k^{2n}$

$$\rho^n = (-1)^n 2^{1-n} \sin(n\varphi), \quad \varphi \in [0, 2\pi). \tag{10}$$

Without lost of generality, from now on we assume that polar equation of  $k^{2n}$  is given by expression:

$$\rho = \sqrt[n]{\sin(n\varphi)}, \quad \varphi \in [0, 2\pi). \tag{11}$$

Let us now consider a congruence  $C(p)$  that consists of circles passing through two given points  $P_{1,2}(0, 0, \pm p)$ , where  $p$  is a positive real number. For the congruence  $C(p)$  and curve  $k^{2n}$  given by (9), a circular surface  $CS(k^{2n}, p)$  is defined as the system of circles from  $C(p)$  that intersect  $k^{2n}$ . According to (CS), if the  $n$ -fold point of  $k^{2n}$  coincide with  $P_1$ , the obtained surface  $CS(k^{2n}, p)$  is entirely spherical surface of order  $2n$  having two real  $n$ -fold points in  $P_1$  and  $P_2$ .




Figure 6

Here we offer a simpler construction of such surface. Let  $k^{2n}$  be a curve in  $xy$ -plane having an  $n$ -fold point in the origin  $O$ , and let  $P \neq O$  be a point on the axis  $z$ . In a plane  $\pi(\varphi)$ , such that  $z \subset \pi$  and  $\varphi = \angle(\pi, x)$ , a circle  $c$  of radius  $PT$ , where  $T = \pi \cap k^{2n}$ , is considered. See Figure 6.

If  $n$  is odd, there is a unique circle  $c$  in every plane  $\pi(\varphi)$ . If  $n$  is even, there could be two or none such circles. For given  $n$  and  $p$  all circles  $c$  determine observed surface  $CS(k^{2n}, \frac{p}{2})$ .

In the plane  $\pi(\varphi)$ , with coordinates  $(\rho, z)$ , the circle  $c$  has an equation

$$(\rho - \rho_S)^2 + (z - z_S)^2 = r^2, \tag{12}$$

where  $\rho_S = \frac{\rho_T}{2}$ ,  $z_S = \frac{p}{2}$ ,  $r^2 = \frac{\rho_T^2}{4} + \frac{p^2}{4}$ , and  $\rho_T$  is given by (11).

Therefore, for  $\varphi \in [0, 2\pi)$ , the equation (12) takes the form

$$\rho^2 - \rho^n \sqrt{\sin n\varphi} + z^2 - pz = 0 \tag{13}$$

which presents the equation of the surface  $CS(k^{2n}, \frac{p}{2})$  in cylindrical coordinates  $(\rho, \varphi, z)$ . If we raise it to the  $n$ -th power, the equation of the surface can be written as

$$((\rho^2 + z^2) + (-pz))^n = \rho^n \sin n\varphi. \tag{14}$$

Since the correspondence between cylindrical and Cartesian coordinates is given by  $\rho = \sqrt{x^2 + y^2}$  and  $\sin \varphi = \frac{y}{\sqrt{x^2 + y^2}}$ , if we use introduced notation  $A_2 = x^2 + y^2 + z^2$  and multiple-angle formulas [25]

$$\sin n\varphi = \begin{cases} (-1)^{(n-1)/2} T_n(\sin \varphi), & n \text{ odd,} \\ (-1)^{n/2-1} \cos \varphi U_{n-1}(\sin \varphi), & n \text{ even,} \end{cases}$$

where  $T_n$  and  $U_n$  are Chebyshev polynomials of the first and second kind, we can write equation (14) in the following form:

$$A_2^n + \sum_{j=1}^{n-1} \binom{n}{j} (-pz)^j A_2^{n-j} = \mathcal{G}^n(x, y) - (-pz)^n \tag{15}$$

where

$$\mathcal{G}^n(x, y) = \begin{cases} (-1)^{\frac{n-1}{2}} (\sqrt{x^2 + y^2})^n T_n\left(\frac{y}{\sqrt{x^2 + y^2}}\right), & n \text{ odd,} \\ (-1)^{\frac{n}{2}-1} x (\sqrt{x^2 + y^2})^{n-1} U_{n-1}\left(\frac{y}{\sqrt{x^2 + y^2}}\right), & n \text{ even.} \end{cases}$$

The properties of the polynomials  $T_n$  and  $U_{n-1}$  ([26], [27]), with  $x^2 + y^2 > 0$ , lead us to the conclusion that  $\mathcal{G}^n$  are homogeneous polynomials of degree  $n$  in  $x$  and  $y$ . Therefore, the equation of constructed surface can be given in the following form:

$$A_2^n + \sum_{j=1}^{n-1} \binom{n}{j} (-pz)^j A_2^{n-j} - \mathcal{T}^n(x, y, z) = 0, \tag{16}$$

where  $\mathcal{T}^n(x, y, z) = \mathcal{G}^n(x, y) - (-pz)^n$  is a homogeneous polynomial of degree  $n$  in  $x$ ,  $y$  and  $z$  determining the equation of tangent cone of the surface at its  $n$ -fold point  $O$ . By using *Mathematica* functions *ChebyshevT* and *ChebyshevU* it is easy to obtain  $\mathcal{T}^n(x, y, z)$  for every  $n$  and  $p$ .

The surface is symmetrical with respect to the plane  $z = \frac{p}{2}$ . Thus, the tangent cone at the  $n$ -fold point  $P$  is symmetrical to the tangent cone at the point  $O$  with respect to the same plane. Some examples of these surfaces are depicted in Figures 7 and 8.




Figure 7: Three examples of surfaces given by equation (15) for  $n = 2, 4, 6, p = 2$  and their  $n^{\text{th}}$  degree tangent cones at the origin. The equations of tangent cones are:  $xy - 2z^2 = 0$  (case a),  $x^3y - xy^3 - 4z^4 = 0$  (case b) and  $3x^5y - 10x^3y^3 + 3xy^5 - 32z^6 = 0$  (case c).




Figure 8: Three examples of surfaces given by equation (15) for  $n = 3, 5, 7, p = 2$  and their  $n^{\text{th}}$  degree tangent cones at the origin. The equations of tangent cones are:  $x^2y - y^3 + 8z^3 = 0$  (case a),  $x^4y - 10x^2y^3 + y^5 + 32z^5 = 0$  (case b) and  $x^6y - 35x^4y^3 + 21x^2y^5 - y^7 + 128z^7 = 0$  (case c).

## References

- [1] A. B. Basset, *An elementary treatise on cubic and quartic curves*, Cambridge, Deighton Bell and Co., London George Bell and Sons, 1901.
- [2] L. P. Eisenhart, *A treatise on the differential geometry of curves and surfaces*, Ginn and Company, Boston - New York - Chicago - London, 1909.
- [3] K. Fladt, *Analytische geometrie spezieller ebener Kurven*, Akademische Verlagsgesellschaft, Frankfurt am Main, 1962.
- [4] A. Goetz, *Introduction to differential geometry*, Addison Wesley Publishing Com., London, 1970.

- [5] S. Gorjanc, *Quartics in  $E^3$  which have a triple point and touch the plane at Infinity through the absolute conic*, Math. Commun. **9** (2004), 67–78.
- [6] S. Gorjanc, *Rose surfaces and their visualizations*, J. Geom. Graphics **13**(1) (2010), 59–67.
- [7] S. Gorjanc, E. Jurkin, *Circular surfaces  $CS(\alpha, p)$* , Filomat **29**(4) (2015), 725–737.
- [8] S. Gorjanc, E. Jurkin, *Generalised Rose surfaces and their visualizations*, Novi Sad J. Math. **45**(2) (2015), 173–185.
- [9] S. Gorjanc, E. Jurkin, *On the special surfaces through the absolute conic with a singular point of the highest order*, Proceedings of the 16th International Conference on Geometry and Graphics, ed. H. P. Schröcker, M. Husty, Innsbruck university press, (2014) ,1168-1173.
- [10] J. Harris, *Algebraic geometry*, Springer, New York, 1995.
- [11] D. Hilbert, S. Cohn-Vossen, *Geometry and the imagination*, Chelsea Publishing Company, New York, 2nd edition, 1990.
- [12] P. H. Johansen, M. Loberg, R. Piene, *Monoid hypersurfaces, geometric modeling and algebraic geometry*, B. Jüttler, R. Piene (ed.), 2008, 55–77.
- [13] E. Kranjčević, *Nožišne krivulje i plohe*, Master Thesis, Faculty of Science, Zagreb, 1967.
- [14] H. Pottmann, L. Shi, M. Skopenkov, *Darboux cyclides and web from circles*, Comp. Aided Geom. Design **29**(1) (2012), 77–97. <http://arxiv.org/pdf/1106.1354v1.pdf>
- [15] K. Rohn, *Ueber die Flächen vierten ordnung mit dreifachen punkte*, Math. Ann. Band **XXIV**, 55–152, B. G. Teubner, Leipzig, 1884.
- [16] G. Salmon, *A treatise on the analytic geometry of three dimensions*, Vol.I., Chelsea Publishing Company, New York, (reprint), 1958.
- [17] G. Salmon, *A treatise on the analytic geometry of three dimensions*, Vol.II., Chelsea Publishing Company, New York, (reprint), 1965.
- [18] G. Salmon, *Higher plane curves*, Chelsea Publishing Company, New York, (reprint), 1960.
- [19] A. A. Savelov, *Ravninske krivulje*, Školska knjiga, Zagreb, 1979.
- [20] H. Wieleitner, *Theorie der ebenen algebraischen Kurven höherer ordnung*, G. J. Göschen'sche Verlagshandlung, Leipzig, 1905.
- [21] *Circular algebraic curve*, (January 6, 2020). In *Wikipedia, The Free Encyclopedia*. Retrieved 16:30, March 5, 2020, from [https://en.wikipedia.org/wiki/Circular\\_algebraic\\_curve](https://en.wikipedia.org/wiki/Circular_algebraic_curve)
- [22] E. W. Weisstein, *Algebraic curve*, From *MathWorld*—A Wolfram Web Resource. <http://mathworld.wolfram.com/AlgebraicCurve.html>
- [23] E. W. Weisstein, *Cyclide*, From *MathWorld*—A Wolfram Web Resource. <http://mathworld.wolfram.com/Cyclide.html>
- [24] E. W. Weisstein, *Partition function P*, From *MathWorld*—A Wolfram Web Resource. <http://mathworld.wolfram.com/PartitionFunctionP.html>
- [25] E. W. Weisstein, *Multiple-angle formulas*, From *MathWorld*—A Wolfram Web Resource. <https://mathworld.wolfram.com/Multiple-AngleFormulas.html>
- [26] E. W. Weisstein, *Chebyshev Polynomial of the First Kind*, From *MathWorld*—A Wolfram Web Resource. <https://mathworld.wolfram.com/ChebyshevPolynomialoftheFirstKind.html>
- [27] E. W. Weisstein, *Chebyshev Polynomial of the Second Kind*, From *MathWorld*—A Wolfram Web Resource. <https://mathworld.wolfram.com/ChebyshevPolynomialoftheSecondKind.html>